

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 057 891 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
06.12.2000 Bulletin 2000/49

(51) Int Cl.7: **C12N 15/29, C12N 15/82,
C12N 5/10, C07K 14/415,
A01H 5/00, A01H 5/10**

(21) Application number: **99201745.9**

(22) Date of filing: **02.06.1999**

(84) Designated Contracting States:

**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**

Designated Extension States:

AL LT LV MK RO SI

(71) Applicants:

- **CENTRUM VOOR PLANTENVEREDELINGS- EN
REPRODUKTIEONDERZOEK
6708 PB Wageningen (NL)**
- **Her Majesty in Right of Canada, represented by
The Minister of Agriculture and Agri-Food
Canada
Ottawa, Ontario K1A 0C6 (CA)**

- **Custers, Jan
6703 CN Wageningen (NL)**
- **Miki, Brian
Ottawa, Ontario K1H 5V1 (CA)**
- **Ouellet, Therese
Nepean, Ontario K2J 3X7 (CA)**
- **Hattori, Jiro
Ottawa, Ontario K1G 1M5 (CA)**
- **Boutillier, Kim
6702 CP Wageningen (NL)**

(74) Representative: **De Hoop, Eric
Octrooibureau Vriesendorp & Gaade
P.O. Box 266
2501 AW Den Haag (NL)**

(72) Inventors:

- **Van Lookeren Campagne, Michiel
6703 CN Wageningen (NL)**

(54) **Use of the BNM3 transcriptional activator to control plant embryogenesis and regeneration processes**

(57) The present invention provides for a gene obtained during the induction of microspore embryogenesis. The protein encoded by this gene renders plant cells

embryogenic, and increases the regenerative capacity of the plant cell. Also disclosed is the regulatory region of this gene and its use for directing the expression of a gene of interest within a suitable host cell.

EP 1 057 891 A1

Description

[0001] The present invention relates to asexual embryo formation and regeneration in plants. More specifically, it relates to processes for producing asexually-derived embryos, and for enhancing regeneration capacity in plants. The present invention also relates to heterologous protein production systems in plants, and the uses thereof.

BACKGROUND OF THE INVENTION

[0002] A typical angiosperm seed consists of three major components, the embryo, the endosperm and the maternal seed coat. Seed development begins with a double fertilization event, in which one sperm cell nucleus fuses with the egg cell nucleus to form the embryo, and a second sperm cell nucleus fuses with two central cell nuclei to form the endosperm. Embryo development itself can be separated into three developmental phases. The first phase of embryo development is one of cell division and morphogenesis, which serves to establish the major tissue types and organ systems of the mature plant. The second phase encompasses a period of rapid cell expansion and is characterized by the synthesis of storage reserves that sustain the embryo during germination and early seedling development. In the final phase of embryo development, the embryo becomes desiccated and enters into a period of developmental arrest or dormancy. All of the above events normally take place while the seed remains attached to the maternal plant.

[0003] Many plant species are capable of producing embryos in the absence of fertilization. This process of asexual embryo development may occur naturally, for example on the leaf margins of *Bryophyllum* (Yarborough, 1923) and *Malaxis* (Taylor, 1967), or within the ovule of apomictic plants (Koltunow, 1995). Apomixis refers to the production of a seed from the maternal ovule tissues in the absence of egg cell fertilization. Asexual embryo development may also be induced *in vitro* from gametophytic or somatic tissue (Mordhorst *et al.*, 1997) or, as shown recently, may be induced by genetic modification of gene expression (Ogas *et al.*, 1997; Lotan *et al.*, 1998).

[0004] Three major mechanisms of apomixis, diplospory, apospory and adventitious embryony, have been observed. Each mechanism differs with respect to the source of the cell that gives rise to the embryo and with respect to the time during ovule development at which the apomictic process is initiated. Diplospory and apospory are considered gametophytic forms of apomixis as they involve the formation of diploid embryo sacs. Adventitious embryony does not involve the production of a mitotically-derived embryo sac.

[0005] In diplospory, the megaspore mother cell does not undergo normal meiosis, but rather divides mitotically to produce a diploid embryo sac instead of the normal haploid embryo sac. One of the cells of the embryo sac functions as the egg cell and divides parthenogenetically (without fertilization) to form an embryo. In some species the unreduced polar nuclei of the embryo sac may fuse to form the endosperm (autonomous endosperm production), the nutritive tissue of the seed, while in other species pollination is necessary for endosperm production (pseudogamy).

[0006] In aposporous apomicts, parthenogenic embryos are produced from additional cells, the aposporous initials, that differentiate from the nucellus. As with the megagametophyte of diplosporous species, the aposporous initial undergoes mitotic divisions to produce a diploid embryo sac. Aposporous embryos are not derived from the megagametophyte and can therefore co-exist within a single ovule with sexually-derived embryos. Autonomous production of endosperm is rare in aposporous species. Aposporous apomicts therefore depend on fertilization of the polar nuclei of a meiotically-derived embryo sac for the production of endosperm.

[0007] With adventitious embryony, embryos are formed directly from sporophytic ovule tissue, such as the integuments or nucellus, via parthenogenesis. Seeds derived from species exhibiting adventitious embryony generally contain multiple asexually-derived embryos and may also contain a single sexually-derived embryo. Plants exhibiting adventitious embryo also rely on the presence of a meiotically-derived embryo sac within the same ovule for endosperm formation.

[0008] In most plant species, the apomictic trait appears to be under the control of a single dominant locus. This locus may encode one or more developmental regulators, such as transcription factors, that in sexually reproducing plants function to initiate gene expression cascades leading to embryo sac and/or embryogenesis, but which are heterochronically or ectopically expressed in apomictic plants (Peacock, 1992; Koltunow, 1993; Koltunow *et al.*, 1995).

[0009] Apomixis is a valuable trait for crop improvement since apomictic seeds give rise to clonal offspring and can therefore be used to genetically fix hybrid lines. The production of hybrid seed is a labour intensive and costly procedure as it involves maintaining populations of genetically pure parental lines, the use of separate pollen donor and male-sterile lines, and line isolation. Production of seed through apomixis avoids these problems in that once a hybrid has been produced, it can be maintained clonally, thereby eliminating the need to maintain and cross separate parental lines. The use of apomictic seed also provides a more cost effective method of multiplying vegetatively-propagated crops, as it eliminates the use of cuttings or tissue culture techniques to propagate lines, reduces the spread of diseases which are easily transmitted through vegetatively-propagated tissues, and in many species reduces the size of the propagule leading to lower shipping and planting costs.

[0010] Although apomixis occurs in a wide range of plant species, few crop species are apomictic. Attempts to in-

produce apomictic traits into crop species by introgression from wild relatives (Ozias-Akins, *et al.*, 1993; WO 97/10704; WO 97/11167) or through crosses between related, but developmentally divergent sexual species (WO 98/33374), have not yielded marketable products. Other approaches have focused on the identification of gene sequences that may be used to identify or manipulate apomictic processes (WO 97/43427; WO 98/36090), however these approaches have not led to methods for the routine production of apomictic plants.

[0011] Mutagenesis approaches have also been attempted to convert sexually reproducing plants such as *Arabidopsis thaliana* (arabidopsis) into apomictic plants (Peacock *et al.*, 1995). For example, a number of recessive "fertilization-independent seed" (*fis*) mutants have been identified that initiate partial embryo and/or endosperm at a low frequency in the absence of fertilization (Chaudhury *et al.*, 1997). However, a number of additional parameters need to be modified in order to obtain true diploid apomictic seed using *fis* mutants.

[0012] Asexually-derived embryos can be induced to form in culture from many gametophytic and somatic plant tissues (Yeung, 1995). Somatic embryos can be obtained from culture of somatic tissues by treating them with plant growth regulators, such as auxins, or auxins in combination with cytokinins. Embryos can also be induced to form in culture from the gametophytic tissues of the ovule (gynogenesis) and the anther (androgenesis, pollen or microspore embryogenesis), either by the addition of plant growth regulators or by a simple stress treatment.

[0013] Several mutants have been identified that may be used to induce efficient production of embryos *in vitro*. These include recessive arabidopsis mutants with altered shoot meristems, for example *primordia timing* (*pt*), *clavata* (*clv*)1 and *clv*3, which were shown to enhance embryogenic callus formation when seedlings were germinated in the presence of auxin (Mordhorst *et al.*, 1998). The altered expression of two arabidopsis genes, *LEAFY COTYLEDON* (*LEC1*; WO 98/37184, Lotan *et al.*, 1998) and *pickle*, have been shown to promote the production of somatic embryos in the absence of added growth regulators. The *LEC1* gene encodes a homologue of the HAP3 subunit of a CCAAT box-binding transcription factor (CBF). The *LEC1* gene controls many aspects of zygotic embryo development including desiccation tolerance and cotyledon identity. Ectopic over-expression of the *LEC1* gene in a *lec1* mutant background results in the production of 2 transgenic lines that occasionally form embryo-like structures on leaves.

[0014] These embryo-like structures express genes, such as those encoding seed storage proteins and oil body proteins, which are normally preferentially expressed in developing embryos. Plants containing a recessive mutant *PICKLE* gene produce a thickened, primary root meristem. Mutant *pickle* roots produce embryo-forming callus when the root tissue is separated from the rest of the plant and placed on minimal medium without growth regulators (Ogas *et al.*, 1997). Mutant *pickle* roots show morphological characteristics of developing seeds, such as oil bodies and, as with *LEC1* over-expressers, accumulate genes preferentially expressed in developing seeds.

[0015] Efficient production of apomictic seed is only likely to be realised through the identification and subsequent modification of developmental regulators, such as transcription factors, that are known to activate gene expression cascades leading to embryogenesis in both sexually-reproducing and apomictic plants. The present invention addresses this need by providing methods for the production of apomictic seeds comprising ectopic over-expression of an embryo-expressed AP2 domain containing transcription factor, BNM3.

SUMMARY OF THE INVENTION

[0016] The present invention relates to asexual embryo formation and regeneration in plants. More specifically, it relates to processes for producing asexually-derived embryos, and for enhancing regeneration capacity in plants.

[0017] According to the present invention there is provided an isolated DNA molecule comprising a nucleotide sequence that hybridizes to SEQ ID NO:5 under stringent conditions, that comprises at least 23 contiguous nucleotides of SEQ ID NO:5, or that is at least 70% homologous with the nucleotide sequence defined by SEQ ID NO:5.

[0018] This invention further relates to an isolated DNA molecule that hybridizes to SEQ ID NO:5 under stringent conditions, comprising a nucleic acid sequence encoding a protein, wherein the protein when present at a sufficient level within a plant cell renders the cell embryogenic, increases the regenerative capacity of the plant cell, or both renders the cell embryogenic and increases the regenerative capacity of the plant cell. Included within the present invention is the above isolated DNA molecule comprising a nucleotide sequence that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5 under stringent conditions, or to the nucleotide sequence as defined within SEQ ID NO:1 or 3 under stringent conditions. Also included within the present invention is a vector comprising the isolated DNA molecule as defined above, wherein the isolated DNA molecule is under control of a regulatory element that directs expression of said DNA in a plant cell. The regulatory element may be a constitutive, inducible, tissue specific or a developmental active, regulatory element.

[0019] This invention also embraces a transformed plant cell, a transformed plant, or seed obtained from a transformed plant, each comprising the vector as defined above.

[0020] This invention relates to an isolated protein encoded by an isolated DNA molecule that hybridizes to the nucleotide sequence defined by SEQ ID NO:5 under stringent conditions, wherein the protein, when present at a sufficient level within a plant cell renders the cell embryogenic, or increases the regenerative capacity of the plant cell.

Also included is a protein encoded by an isolated DNA molecule that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5 under stringent conditions, or to the nucleotide sequence as defined within SEQ ID NO:1 or 3 under stringent conditions. This invention also embraces an isolated DNA molecule that encodes a protein as defined by SEQ ID NO:2 or SEQ ID NO:4. The invention also pertains to a protein comprising at least 70% homology with the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:4, or comprises from about 30 to about 541 amino acids of the sequence disclosed in SEQ ID NO:2, or comprises from about 30 to about 561 amino acids of the sequence disclosed in SEQ ID NO:4.

[0021] The present invention is also directed to a method of producing asexually derived embryos comprising:

- i) transforming a plant cell with a vector comprising an isolated DNA molecule that hybridizes to SEQ ID NO:5 under stringent conditions and which encodes a protein that when present at a sufficient level within the plant cell renders the plant cell embryogenic, or increases the regenerative capacity of the plant cell, or that comprises a nucleotide sequence that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5 under stringent conditions, or to the nucleotide sequence as defined within SEQ ID NO:1 or 3 under stringent conditions;
- ii) growing the plant cell to produce transformed tissue;
- iii) selecting the transformed tissue for occurrence of the isolated DNA molecule; and
- iv) assaying the transformed tissue for asexual embryo formation.

[0022] This invention also relates to the above method where the step of assaying (step iv)) involves assaying for somatic embryos, gametophytically-derived embryos, adventitious embryony, diplospory, or for haploid parthenogenesis of the embryo sac.

[0023] The present invention also embraces a method of producing an apomictic plant comprising:

- i) transforming a plant cell with a vector comprising an isolated DNA molecule that hybridizes to SEQ ID NO:5 under stringent conditions and which encodes a protein that when present at a sufficient level within said plant cell renders the plant cell embryogenic, or increases the regenerative capacity of the plant cell, or that comprises a nucleotide sequence that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5 under stringent conditions, or to the nucleotide sequence as defined within SEQ ID NO:1 or 3 under stringent conditions;
- ii) selecting the transformed plant for occurrence of the isolated DNA molecule; and
- iii) assaying the transformed plant for asexual embryo formation.

[0024] This invention also relates to the above method where the step of assaying (step iii)) involves assaying for asexually-derived embryos, somatic embryos, gametophytically-derived embryos, adventitious embryony, diplospory, or for haploid parthenogenesis of the embryo sac.

[0025] The present invention is also directed to a method of producing asexually derived embryos comprising:

- i) transiently transforming a plant cell with a vector comprising an isolated DNA molecule that hybridizes to SEQ ID NO:5 under stringent conditions and which encodes a protein that when present at a sufficient level within the plant cell renders the plant cell embryogenic, or increases the regenerative capacity of said plant cell, or that comprises a nucleotide sequence that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5 under stringent conditions, or to the nucleotide sequence as defined within SEQ ID NO:1 or 3 under stringent conditions;
- ii) growing the transiently transformed plant cell to produce transiently transformed tissue;
- iii) assaying the transiently transformed tissue for asexual embryo formation

[0026] This invention is directed to the above method where the step of assaying (step iii)) involves assaying for asexually-derived embryos, somatic embryos, gametophytically-derived embryos, adventitious embryony, diplospory, or for haploid parthenogenesis of the embryo sac.

[0027] The present invention also presents a method of modifying the regenerative capacity of a plant comprising

- i) transforming a plant cell with a vector comprising an isolated DNA molecule that hybridizes to SEQ ID NO:5 under stringent conditions and which encodes a protein that when present at a sufficient level within the plant cell renders the plant cell embryogenic, or increases the regenerative capacity of said plant cell, or that comprises a nucleotide sequence that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5 under stringent conditions, or to the nucleotide sequence as defined within SEQ ID NO:1 or 3 under stringent conditions;
- ii) growing the transformed plant cell to produce transformed tissue; and
- iii) assaying the transformed tissue for enhanced regeneration as compared to wild type tissue.

[0028] This invention also embraces the above method wherein step iii) includes assaying in the absence of a growth

regulator

[0029] The present invention also relates to a method of modifying the regenerative capacity of a plant comprising:

- i) transiently transforming a plant cell with a vector comprising an isolated DNA molecule that hybridizes to SEQ ID NO:5 under stringent conditions and which encodes a protein that when present at a sufficient level within the plant cell renders the plant cell embryogenic, or increases the regenerative capacity of the plant cell, or that comprises a nucleotide sequence that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5 under stringent conditions, or to the nucleotide sequence as defined within SEQ ID NO:1 or 3 under stringent conditions;
- ii) growing the transiently transformed plant cell to produce transiently transformed tissue;
- iii) assaying the transformed tissue for enhanced regeneration as compared to wild type tissue.

[0030] This invention also embraces the above method wherein step iii) includes assaying in the absence of a growth regulator

[0031] The present invention also relates to a method of selecting a transformed plant comprising:

- i) transforming a normally non-regenerative plant with a vector comprising an isolated DNA molecule that hybridizes to SEQ ID NO:5 under stringent conditions and which encodes a protein that when present at a sufficient level within the plant cell renders the plant cell embryogenic, or increases the regenerative capacity of said plant cell, or that comprises a nucleotide sequence that hybridizes to nucleotides 1620- 4873 of SEQ ID NO:5 under stringent conditions, or to the nucleotide sequence as defined within SEQ ID NO:1 or 3 under stringent conditions; and
- ii) determining whether the transformed plant is able to regenerate under conditions in which the normally non-regenerative plant does not regenerate.

[0032] The present invention is also directed to an isolated DNA molecule comprising a DNA sequence that hybridizes to nucleotides 1-1619 of SEQ ID NO:5 under stringent conditions, or that comprises at least 22 contiguous nucleotides within nucleotides 1-1619 of SEQ ID NO:5. Also included within the scope of the present invention is a vector comprising the isolated DNA molecule as just defined, operably associated with a gene of interest, wherein the isolated DNA molecule directs the expression of the gene of interest within a plant cell. The gene of interest may be heterologous with respect to the isolated DNA molecule. The gene of interest may be selected from the group consisting of a pharmaceutically active protein, antibody, industrial enzyme protein supplement, nutraceutical, storage protein, animal feed and animal feed supplement. This invention also includes a transformed plant cell, a transformed plant, or seed obtained from the transformed plant, comprising the vector as just defined.

[0033] Furthermore, the present invention includes a method for directing the expression of a gene of interest within a developing embryo of a plant comprising transforming said plant with a vector containing an isolated DNA molecule that hybridizes to nucleotides 1-1619 of SEQ ID NO:5 under stringent conditions, or that comprises at least 22 contiguous nucleotides within nucleotides 1-1619 of SEQ ID NO:5.

[0034] This invention also pertains to a method of producing a protein of interest comprising

- i) transforming a plant with at least one vector, comprising an isolated DNA molecule that hybridizes to SEQ ID NO:5 under stringent conditions and which encodes a protein that when present at a sufficient level within the plant cell renders the plant cell embryogenic, or increases the regenerative capacity of said plant cell, or that comprises a nucleotide sequence that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5 under stringent conditions, or to the nucleotide sequence as defined within SEQ ID NO:1 or 3 under stringent conditions to produce a transformed plant;
- ii) selecting the transformed plant for occurrence of the isolated DNA molecule; and
- iv) growing the transformed plant in order to produce the protein of interest, wherein expression of the protein of interest is induced by the expression product of said isolated DNA.

[0035] This method may also comprise transforming the plant with a second vector comprising a nucleotide sequence encoding the protein of interest under the control of a regulatory element, wherein the regulatory element induced by the expression product of the isolated DNA. Furthermore, this method may also be used to produce a protein of interest wherein the protein of interest is a native protein.

BRIEF DESCRIPTION OF THE DRAWINGS

[0036] These and other features of the invention will become more apparent from the following description in which reference is made to the appended drawings wherein:

Figure 1 shows a schematic representation of the effect of culture temperature on the developmental fate of isolated microspores and pollen of *Brassica napus*. Late uninucleate microspores and early binucleate pollen cultured at 25 ° C or lower continue to divide and form functional pollen grains (gametophytic), while the same microspores and pollen cultured at 32 ° C undergo numerous sporophytic divisions, leading to the formation of haploid embryos (embryogenic). Late uninucleate microspores and early binucleate pollen cultured for one day at 25 ° C, followed by culture at 32 ° C may undergo gametophytic divisions, but form neither embryos nor mature pollen grains (non-embryogenic).

Figure 2 shows the alignment of the DNA sequences depicted in SEQ ID NO:1 and SEQ ID NO:3. The ATG and TAG translation initiation and translation termination codons are shown in bold. Identical nucleotides are indicated by (*) and gaps are indicated by (-).

Figure 3 shows the alignment of the predicted protein sequences encoded by the DNA of SEQ ID NO:1 and SEQ ID NO:3. The amino acid sequence of the first AP2 domain repeat (repeat 1) and the second AP2 domain repeat (repeat 2), are shown in bold. Identical amino acids are indicated by an asterisk (*) and mismatches by a dot (.) below the sequence alignment.

Figure 4 shows the presence of two *BNM3* genes in the *Brassica napus* genome. A DNA gel blot containing restriction digests of *B. napus* c.v. Topas genomic DNA was hybridized to a *BNM3A* cDNA fragment under high stringency conditions. The *BNM3A* cDNA hybridizes to two DNA fragments under these conditions. These fragments correspond to the *BNM3A* and *BNM3B* genes. The position of the molecular size markers (Lambda DNA *Hind* III restriction fragments) is indicated to the left of the figure. The restriction enzymes used to digest the DNA are indicated above the blot.

Figure 5 shows the alignment of the predicted protein sequence encoded by the DNA of SEQ ID NO:1 (*BNM3A*) with the predicted protein sequences of other AP2 domain proteins. The amino acid sequence of *BNM3A*, beginning at position 208, and spanning the first AP2 domain repeat (AP2 domain repeat 1), the second AP2 domain repeat (AP2 domain repeat 2), and the linker region lying between the two repeats (linker), was aligned with the amino acid sequence of other proteins containing two AP2 domains. The amino acid similarity in this region ranges from 53% for APETALA2 to 80% for ZMMHCF1. Identical amino acids are indicated by (*) and gaps are indicated by (-). Protein names are indicated on the left and are abbreviated as follows: ANT, AINTEGUMENTA (accession number U41339); ZM, ZMMHCF1 (accession number Z47554); GL15, GLOSSY15 (accession number U41466); AP2, APETALA2 (accession number U12546).

Figure 6 shows the results of gel blot analysis with a *BNM3A* cDNA fragment performed on RNA extracted from the indicated tissues. RNA gel blots contain either 5 µg (a) or 20 µg (b, c) of total RNA. Figure 6A shows the pattern of *BNM3* expression in microspore embryo cultures. RNA was isolated from late uninucleate microspores and early binucleate pollen at the time of collection (pollen 0d), after four days in culture at 32 ° C (+ embryo), after four days in culture at 25 ° C (pollen 4d), after one day of culture at 25 ° C, followed by three days of culture at 32 ° C (- embryo) and microspore-derived embryos at the globular, heart, torpedo, 21 day old cotyledon (21 d cot), 28 day old cotyledon (28 d cot) and 42 day old cotyledon (42 d cot) stage of development. *BNM3* expression is detected in embryogenic microspores and developing microspore-derived embryos, but is absent from developing microspores and pollen collected prior to tissue culture and in non-embryogenic samples. The exposure time was seven days. Figure 6B shows that *BNM3* gene expression is detected in developing seeds. Seeds were collected at various days after pollination (DAP). These points in development correspond approximately to the globular (7 d), heart (14 d), torpedo (18 d), early cotyledon (21 d), mid cotyledon (28 d, 35 d) and late cotyledon (42 d) stages of development. The exposure time was 14 days. Figure 6C shows that *BNM3* gene expression is not detected in non-seed tissues. Roots and leaves were collected from 14 day old greenhouse grown plants. Entire flowers as well as excised anthers and pistils were collected from opened flower buds just prior to anthesis. Small and large buds refer to closed flower buds of less than 5 mm or greater than 5 mm in length, respectively. Siliques were collected 16 days after pollination. The exposure time was 14 days.

Figure 7 shows the phenotype of *Brassica napus* and arabidopsis plants transformed with constructs containing the *BNM3* gene under control of a modified *POLYUBIQUITIN* promoter (B) and double enhanced 35S promoter containing an AMV translational enhancer (A, C-E). Figure 7A shows embryo structures on the leaf margin of a *Brassica* T1 seedling. Figure 7B shows embryo structures on the petiole of an arabidopsis T2 seedling. Figure 7C shows embryo structures on the cotyledon of an arabidopsis T1 seedling. Figure

7D shows a scanning electron micrograph of the abaxial side of an arabidopsis T1 cotyledon. Note the bipolar nature of the embryos, as well as the emergence of a secondary embryo from the surface of a primary embryo (asterisk). Figure 7E shows a semi-thin section through one of the cotyledons of the T1 seedling shown in (Figure 7C). Note the presence of all the major organs and tissue elements of embryo, as well as the development of new embryos on the flanks of the shoot apical meristems and the cotyledons.

Figure 8 shows the increased regenerative capacity of arabidopsis plants transformed with a construct containing the *BNM3B* gene under control of a modified *POLYUBIQUITIN* promoter. Figure 8A shows wild-type and transgenic leaf and hypocotyl explants on medium containing growth regulators. Figure 8B shows wild-type and transgenic roots on medium containing growth regulators. Figure 8C shows wild-type and transgenic leaf and hypocotyl explants on medium without growth regulators. Figure 8D shows wild-type and transgenic root explants on medium without growth regulators.

DESCRIPTION OF PREFERRED EMBODIMENT

[0037] The present invention relates to asexual embryo formation and regeneration in plants. More specifically, it relates to processes for producing asexually-derived embryos, and for enhancing regeneration capacity in plants. The present invention also relates to heterologous protein production systems in plants, and the uses thereof.

[0038] Genes preferentially expressed during the induction of *Brassica napus* c.v. Topas microspore embryogenesis were isolated via subtractive screening. Seven independent cDNA clones, comprising six unique DNA sequences were found to be differentially expressed between cDNA libraries prepared from embryogenic and non-embryogenic microspore cultures. Several of these *BNM* (for *Brassica napus* microspore embryo) clones, *BNM3A* (SEQ ID NO:1) and *BNM3B* (SEQ ID NO:3), were characterized as described herein. *BNM3A* and *BNM3B* encode the amino acid sequences disclosed in SEQ ID NO:2, and SEQ ID NO:4, respectively. The genomic sequence of *BNM3A* (SEQ ID NO:5), including the regulatory region (nucleotides 1-1619 of SEQ ID NO:5), was also obtained.

[0039] "Regeneration", as used herein, refers to a morphogenetic response that results in the production of new tissues, organs, embryos, whole plants or fragments of whole plants that are derived from a single cell, or a group of cells. Regeneration may proceed indirectly via a callus phase or directly, without an intervening callus phase. "Regenerative capacity" refers to the ability of a plant cell to undergo regeneration.

[0040] By "embryogenic cell", it is meant a cell that has completed the transition from either a somatic or a gametophytic cell to a state where no further applied stimuli are necessary to produce a somatic or gametophytic embryo, respectively.

[0041] By "regulatory element" it is meant those that include developmentally regulated, tissue specific, inducible and constitutive regulatory elements. A regulatory element that is developmentally regulated, or controls the differential expression of a gene under its control, is activated within certain organs or tissues of an organ at specific times during the development of that organ or tissue. However, some regulatory elements that are developmentally regulated may preferentially be active within certain organs or tissues at specific developmental stages, they may also be active in a developmentally regulated manner, or at a basal level in other organs or tissues within the plant as well, such regulatory elements are considered "tissue specific". Regulatory elements may be found either upstream, within, downstream, or a combination thereof, of the coding region of a gene.

[0042] An inducible regulatory element is one that is capable of directly or indirectly activating transcription of one or more DNA sequences or genes in response to an inducer. In the absence of an inducer the DNA sequences or genes will not be transcribed. Typically the protein factor, that binds specifically to an inducible regulatory element to activate transcription, is present in an inactive form which is then directly or indirectly converted to the active form by the inducer. The inducer can be a chemical agent such as a protein, metabolite, growth regulator, herbicide or phenolic compound or a physiological stress imposed directly by heat, cold, salt, or toxic elements or indirectly through the action of a pathogen or disease agent such as a virus. A plant cell containing an inducible regulatory element may be exposed to an inducer by externally applying the inducer to the cell or plant such as by spraying, watering, heating or similar methods.

[0043] A constitutive regulatory element directs the expression of a gene throughout the various parts of a plant and continuously throughout plant development. Examples of known constitutive regulatory elements include promoters associated with the CaMV 35S transcript (Odell et al., 1985, *Nature*, 313: 810-812), the rice actin 1 (Zhang et al, 1991, *Plant Cell*, 3: 1155-1165) and triosephosphate isomerase 1 (Xu et al, 1994, *Plant Physiol*, 106: 459-467) genes, the maize ubiquitin 1 gene (Corno et al, 1993, *Plant Mol. Biol.* 29: 637-645), the *Arabidopsis* ubiquitin 1 and 6 genes (Holtorf et al, 1995, *Plant Mol. Biol.* 29: 637-646), and the tobacco translational initiation factor 4A gene (Mandel et al, 1995 *Plant Mol. Biol.* 29: 995-1004).

[0044] By "gene of interest" it is meant any gene that is to be expressed in a transformed plant. Such a gene of interest may include, but is not limited to, a gene that encodes a pharmaceutically active protein, for example growth

factors, growth regulators, antibodies, antigens, their derivatives useful for immunization or vaccination and the like. Such proteins include, but are not limited to, interleukins, insulin, G-CSF, GM-CSF, hPG-CSF, M-CSF or combinations thereof, interferons, for example, interferon- α , interferon- β , interferon- γ , blood clotting factors, for example, Factor VIII, Factor IX, or tPA or combinations thereof. A gene of interest may also encode an industrial enzyme, protein supplement, nutraceutical, or a value-added product for feed, food, or both feed and food use. Examples of such proteins include, but are not limited to proteases, oxidases, phytases chitinases, invertases, lipases, cellulases, xylanases, enzymes involved in oil biosynthesis etc. Other protein supplements, nutraceuticals, or a value-added products include native or modified seed storage proteins and the like.

[0045] The present invention is further directed to a chimeric gene construct containing a DNA of interest operatively linked to a regulatory element of the present invention. Any exogenous gene, or gene of interest, can be used and manipulated according to the present invention to result in the expression of the exogenous gene.

[0046] The activation of the expression of a gene of interest may also be under the control of a regulatory element that itself is activated by a BNM3 protein. For example, which is not to be considered limiting, a gene of interest may be fused to the napin promoter, and the napin promoter may be induced by BNM3. Furthermore, a gene of interest may be expressed within somatic tissues under the control of one or more regulatory elements induced by BNM3, so that, as will be described in more detail below, the somatic tissue develops into a seed-like structure comprising embryogenic cells, and these seed-like structures produce the products of the gene of interest.

[0047] The chimeric gene construct of the present invention can further comprise a 3' untranslated region. A 3' untranslated region refers to that portion of a gene comprising a DNA segment that contains a polyadenylation signal and any other regulatory signals capable of effecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by effecting the addition of polyadenylic acid tracks to the 3' end of the mRNA precursor. Polyadenylation signals are commonly recognized by the presence of homology to the canonical form 5' AATAAA-3' although variations are not uncommon. Examples of suitable 3' regions are the 3' transcribed non-translated regions containing a polyadenylation signal of *Agrobacterium* tumor inducing (Ti) plasmid genes, such as the nopaline synthase (*Nos* gene) and plant genes such as the soybean storage protein genes and the small subunit of the ribulose-1, 5-bisphosphate carboxylase (ssRUBISCO) gene. The 3' untranslated region from the structural gene of the present construct can therefore be used to construct chimeric genes for expression in plants.

[0048] The chimeric gene construct of the present invention can also include further enhancers, either translation or transcription enhancers, as may be required. These enhancer regions are well known to persons skilled in the art, and can include the ATG initiation codon and adjacent sequences. The initiation codon must be in phase with the reading frame of the coding sequence to ensure translation of the entire sequence. The translation control signals and initiation codons can be from a variety of origins, both natural and synthetic. Translational initiation regions may be provided from the source of the transcriptional initiation region, or from the structural gene. The sequence can also be derived from the regulatory element selected to express the gene, and can be specifically modified so as to increase translation of the mRNA.

[0049] To aid in identification of transformed plant cells, the constructs of this invention may be further manipulated to include plant selectable markers. Useful selectable markers include enzymes which provide for resistance to an antibiotic such as gentamycin, hygromycin, kanamycin, and the like. Similarly, enzymes providing for production of a compound identifiable by colour change such as *GUS* (β -glucuronidase), fluorescence, or luminescence, such as luciferase are useful.

[0050] Also considered part of this invention are transgenic plants containing a gene or chimeric gene construct of the present invention comprising a *BNM3* gene, a regulatory element obtained from *BNM3*, or the coding region from *BNM3* in operative association with a constitutive, developmental or inducible regulatory element, or a combination thereof. Methods of regenerating whole plants from plant cells are known in the art. In general, transformed plant cells are cultured in an appropriate medium, which may contain selective agents such as antibiotics, where selectable markers are used to facilitate identification of transformed plant cells. Once callus forms, shoot formation can be encouraged by employing the appropriate plant hormones in accordance with known methods and the shoots transferred to rooting medium for regeneration of plants. The plants may then be used to establish repetitive generations, either from seeds or using vegetative propagation techniques. The constructs of the present invention can be introduced into plant cells using Ti plasmids, Ri plasmids, plant virus vectors, direct DNA transformation, micro-injection, electroporation, biolistics etc. For reviews of such techniques see for example Weissbach and Weissbach, *Methods for Plant Molecular Biology*, Academy Press, New York VIII, pp. 421-463 (1988); Geierman and Corey, *Plant Molecular Biology*, 2d Ed. (1988); and Miki and Iyer, *Fundamentals of Gene Transfer in Plants*, In *Plant Metabolism*, 2d Ed. DT. Dennis, DH Turpin, DD Lefebvre DB Layzell (eds), Addison Wesley, Langmans Ltd. London, pp. 561-579 (1997). The present invention further includes a suitable vector comprising the gene or the chimeric gene construct.

[0051] A class of genes have been isolated from *Brassica napus* microspore embryo cultures. These genes have been found to be important regulators of embryogenesis by their ability to induce the formation of asexually-derived embryos when ectopically expressed in the vegetative tissues of plants. These genes are hereinafter indicated as

BNM3 genes (*Brassica napus* microspore embryo). SEQ ID NO. 1 depicts the cDNA of *BNM3A*, SEQ ID NO. 3 depicts the cDNA of *BNM3B*, and the genomic sequence for *BNM3A* is given in SEQ ID NO:5. The promoter of *BNM3A* lies within nucleotides 1-1619 of SEQ ID NO:5. The predicted protein sequences encoded by the DNAs of SEQ ID NO. 1 and 3 are outlined in SEQ ID NOs. 2 and 4, respectively.

[0052] By "*BNM3*" or "*BNM3* gene", it is meant the sequence of oligonucleotides as disclosed in SEQ ID NOs: 1, 3 or 5, or fragments, derivatives, or mutations thereof, or oligonucleotide sequences that exhibit at least 70% homology or similarity, with a fragment or derivative of the sequences disclosed in SEQ ID NOs 1, 3, or the coding region (nucleotides 1620-4873) of SEQ ID NO:5, as determined using oligonucleotide alignment algorithms (for example, but not limited to a BLAST or FASTA). Furthermore, oligonucleotides that associate with these sequences under conditions of high stringency, for example, but not to be limited to, hybridization to gel blots at about 65°C followed by wash conditions at 0.1X SSC, 65°C, are also considered *BNM3* genes. "*BNM3* gene" also includes DNA molecules that comprises at least 23 contiguous nucleotides of SEQ ID NOs:1, 3 or 5, or at least 22 contiguous nucleotides within nucleotides 1-1619 of SEQ ID NO:5. A fragment of *BNM3*, that comprises at least 22 contiguous nucleotides may be used as a probe for the identification of nucleotides related to *BNM3* regulatory, or coding, regions within an organism. Furthermore, molecules comprising at least 23 contiguous nucleotides of SEQ ID NOs: 1, 3 or 5 and that encode a protein, or an active fragment thereof, that when present at a sufficient level within a plant cell renders the cell embryogenic, increases the regenerative capacity of the plant cell, or renders the cell embryogenic and increases the regenerative capacity of the plant cell are also considered to be *BNM3* genes. Preferably, a *BNM3* gene comprises from about 50 to about 1981 nucleotides of SEQ ID NOs: 1 or 3, or about 50 to about 4858 nucleotides from the coding region (1620-4858) of SEQ ID NO:5.

[0053] By "*BNM3* regulatory region" it is meant the sequence of oligonucleotides from 1-1619 in SEQ ID NO:5, or a fragment, derivative, or mutation thereof. Furthermore, a *BNM3* regulatory region also comprises a nucleotide sequence that associates with a sequence from 1-1619 of SEQ ID NO:5 under conditions of high stringency, for example, but not to be limited to, hybridization to gel blots at about 65°C followed by wash conditions at 0.1X SSC, 65°C, or that exhibits at least 70% homology or similarity, with a fragment or derivative of the sequences disclosed in nucleotides 1-1619 of SEQ ID NO:5, as determined using oligonucleotide alignment (for example, but not limited to a BLAST or FASTA search).

[0054] By "*BNM3* protein" it is meant a protein, or a biologically active fragment thereof, that renders a plant cell embryogenic, increases the regenerative capacity of the plant cell, or renders the cell embryogenic, increases the regenerative capacity of the plant cell, and that is encoded by a *BNM3* gene, as defined above. Preferably, a *BNM3* protein comprises from about 30 to about 541 amino acids of the sequence disclosed in SEQ ID NO:2, or from about 30 to about 561 amino acids of the sequence disclosed in SEQ ID NO: 4. However, *BNM3* protein may also be defined as a protein having at least 70% homology with either SEQ ID NO:2 or 4.

[0055] Search of the sequence databases indicated that the *BNM3* translation products contain two copies of an AP2 domain (Figure 3; see also SEQ ID NO: 2 for *BNM3A*, and SEQ ID NO: 4 for *BNM3B*). The AP2 domain was first identified in APETALA2, an arabinidopsis protein that regulates meristem identity, floral organ specification, seedcoat development and floral homeotic gene expression (Jofuku *et al.*, 1994), but has since been identified in a wide range of proteins with diverse functions.

[0056] The AP2 domain is usually between 58 to 68 amino acids in length and contains a conserved central core of 18 amino acids, characterized by its ability to form an amphipathic α helix, a structure thought to mediate protein-protein interactions. The ability of a number of AP2 domain containing proteins to bind DNA, coupled with the presence of putative nuclear localization signals and acidic regions that may function as transcriptional activators suggests these proteins function as transcription factors.

[0057] Two phylogenetically distinct classes of AP2 domain proteins have been identified; proteins with a single AP2 domain (EREBP-like) and proteins with two AP2 domains (AP2-like, (Zhou, 1997)). The proteins encoded by the genes of this invention represent unique members of the latter class of proteins.

[0058] Accordingly, an aspect of the present invention provides for an isolated DNA molecule that comprises a sequence encoding a protein that contains two AP2 domains. The protein, when present at a sufficient level in a plant cell, renders the cell embryogenic, increases the regenerative capacity of the cell, or both renders the cell embryogenic and increases the regenerative capacity of the cell.

[0059] Analysis of *BNM3* expression during microspore-derived embryo development, seed development, or non-seed tissue development, using Northern (Figure 6) indicated that the *BNM3* genes are preferentially expressed in embryogenic microspore cultures, microspore-derived embryos and seeds. *BNM3* transcripts were not detected in any of the non-seed tissues tested.

[0060] *BNM3* mRNA is detected in microspore cultures induced to undergo embryogenesis, as well as in the subsequent globular, heart, torpedo and cotyledon stages of microspore-derived embryo development (e.g. Figure 6A). RNAs are also detected within developing seeds, 14 days after pollination (14 DAP), corresponding to the heart stage of embryo development. *BNM3* expression increases during the early (21 DAP) and mid-cotyledon (28 DAP) stages

of embryo development and remains constant thereafter (Figure 6B).

[0061] Constitutive expression of *BNM3* resulted in the formation of somatic embryos on vegetative structures such as cotyledons, petioles, leaf blades and the shoot apical meristem of plants (Figure 7). In these experiments *BNM3* cDNAs were placed under the control of two separate constitutive promoter constructs, a modified sunflower *POLYU-BIQUITIN* promoter construct, and a double enhanced *35S* promoter construct containing an AMV translational enhancer, however, it is to be understood that any suitable constitutive promoter may be used for this purpose. Such *BNM3*-derived ectopic embryos contain all of the organ systems and tissue layers found in the developing zygotic embryo in that these embryos are bipolar (Figure 7E), consist of an axis, a hypocotyl and radicle region, shoot and root meristems, and cotyledons. In addition, each organ system contained the characteristic radial arrangement of three specialized tissue layers (epidermis, ground parenchyma and provascular tissue) found in zygotic embryos. Continued expression of the *BNM3* gene within the developing ectopic embryo leads to a reiteration of the embryo-forming process, with the result that new embryos are continuously formed on the surface of pre-existing embryos (Figure 7E).

[0062] Constitutive expression of *BNM3* results in the increased ability of a plant to regenerate shoots *in vitro* in the presence of added growth regulators. Root explants from transgenic plants ectopically expressing *BNM3* show at least a 5-fold increase in shoot regeneration in the presence of hormones as compared to root explants obtained from wild-type plants (Figure 8A,B). Shoots also developed faster in the transgenic explants, compared to the wild-type. Wild-type leaf and hypocotyl explants initially responded by producing callus on the cut end of the petiole (Figure 8B) followed by callus formation along the length of the petiole. In contrast, explants from transgenic lines immediately produced new shoots (Figure 8B) or roots from the cut end of the petiole. Explants that initially produced roots eventually also produced shoots.

[0063] Transgenic explants, constitutively expressing *BNM3* were also able to regenerate in the absence of added growth regulators. These explants, when placed on media lacking growth regulators regenerated shoots either from the cut end of the leaf and hypocotyl explants or from the nodule-like structures of root explants (Figures 8C,D). In all cases regenerated shoots developed, rooted, flowered and set seed. Conversely, wild-type leaf and hypocotyl explants placed on medium lacking growth regulators occasionally produce callus or roots at the cut end of the leaf petiole, however no shoots form from these structures (Figure 8C,D).

[0064] It is also considered within the scope of the present invention, that expression of *BNM3* may be used to initiate a developmental cascade within a transformed plant or plant cell. This cascade may arise as a result of the stable integration of a DNA-based vector expressing *BNM3* within a transformed plant, however, such a cascade may also arise as a result of transient expression of *BNM3*, and does not require the stable integration of the *BNM3*-based vector within a plant cell. These transient approaches may be useful for inducing somatic embryogenesis, gametophytically-derived embryogenesis, or increasing the regenerative capacity of a plant or plant cell.

[0065] Plants in which a *BNM3* gene is ectopically expressed exhibit advantageous qualities including:

- formation of asexually derived embryos;
- increased regenerative capacity of tissue explants;
- the ability of tissue explants to regenerate in the absence of added plant growth regulators; and
- the expression of seed components in non-seed organs in which *BNM3* is ectopically expressed.

[0066] Furthermore, plants that ectopically express at least one *BNM3* gene can be used for the production of recombinant proteins using seed specific regulatory elements.

[0067] For the applications of *BNM3* as described below, it will be advantageous to obtain a high level of the *BNM3* transcript and/or *BNM3* protein in order to obtain plants in which the phenotype is highly penetrant. This may be obtained by using genetic elements such as introns, transcriptional enhancers or translational enhancers which are known to enhance gene or protein expression levels.

[0068] The *BNM3* sequences of the present invention may be used for several applications including, but not limited to the control of embryo processes, the control of regeneration processes, the use of regulatory sequences for targeted gene expression, the use of *BNM3* sequences as selectable markers of transformed plants, or for embryogenic cells. These applications are disclosed in more detail below.

Use of *BNM3* Sequences to Control Embryogenic Processes

[0069] As described herein, *BNM3* genes play an important role in initiation and maintenance of embryo development. *BNM3* genes have been found in a wide range of members of the plant kingdom. Regulatory regions obtained from these genes may be used to control the transcription of *BNM3* or a derivative or fragment thereof, or any gene of interest, using methods known to one of skill in the art.

[0070] Ectopic expression of a *BNM3* gene is sufficient to induce recurrent formation of asexually derived embryos

on the vegetative tissues of plants (see example 4). Depending upon the promoter used, ectopic over-expression of *BNM3* genes may be used to produce somatic or gametophytic embryos. Somatic or gametophytic embryos may be obtained by expressing a *BNM3* gene under the control of a constitutive regulatory element, as is shown in Example 5, or may also be obtained by expressing a *BNM3* gene under the control of tissue specific or developmentally regulated elements, inducible elements derived from either plant or non-plant genes or through transient expression. In this respect, chemical induction systems (e.g. see Gatz and Lenk, 1998, which is incorporated by reference) or transient expression using methods which do not result in stable integration of the *BNM3* gene, or which make direct use of the *BNM3* protein e.g. microprojectile bombardment of DNA or protein may also be employed.

[0071] Temporal and/or spatial restriction of *BNM3* expression using inducible, tissue specific or developmentally regulated elements, is preferred when recurrent embryogenesis is not a desirable trait. The regulatory elements used to restrict *BNM3* to a specific developmental stage or cell type will depend on the application. For example, regulatory elements that may be used to express *BNM3* for the production of microspore-derived embryos include, but are not limited to, those of the class I low molecular weight heat shock inducible gene, *GMHSP17.3B* (Zarsky *et al.*, 1995, which is incorporated by reference), or microspore/pollen expressed genes such as *NTM19* (Custers *et al.*, 1997, EP 790,311, which are incorporated by reference), *BCP1* (Xu *et al.*, 1995, which is incorporated by reference), *LAT52* (Twell *et al.*, 1989, which is incorporated by reference), *BNM1* (Treacy *et al.* 1997, which is incorporated by reference) and *APG* (Roberts *et al.*, 1993, which is incorporated by reference).

[0072] Examples of regulatory elements that may be used to express *BNM3* for the production of somatic embryos include, but are not limited to, those of genes activated by plant growth regulators which are routinely used to induce somatic embryogenesis in tissue culture. Specific examples, which are to be considered non-limiting, include the cytokinin inducible *IB6* and *CK11* genes (Brandsstatter and Kieber, 1998; Kakimoto, 1996, which are incorporated by reference) and the auxin inducible element DR5 (Ulmasov *et al.*, 1997, which is incorporated by reference). However, it is to be understood that other regulatory elements may be included for the expression of *BNM3* in plants.

[0073] Furthermore, examples of gene regulatory elements suitable for directing expression of *BNM3* to obtain adventitious embryony include, but are not limited to, those obtained from the ovule and embryo expressed *SERK* gene (Schmidt *et al.*, 1997 which is incorporated by reference), the ovule expressed *AGL11* gene (Roundley *et al.*, 1995, which is incorporated by reference), the nucellus expressed *NUC1* gene (Doan *et al.*, 1996; WO 98/08961, which are incorporated by reference), or the inner integument-expressed genes, *FBP7* (Angenent *et al.*, 1995, which is incorporated by reference) and *SC4* (US application 09/059,909, filed April 13, 1998, which is incorporated by reference) genes.

[0074] According to one aspect of the present invention there is provided a method for the efficient production of microspore-derived embryos in plants. This method involves:

- i) transforming a plant of interest, for example, *Brassica napus* (using transformation techniques known to one of skill, for example, DeBlock *et al.*, 1989, Clough and Bent 1998, Vergunst *et al.* 1998, Klein *et al.* 1987, which are incorporated herein by reference) with a vector construct, or isolated DNA, consisting of a *BNM3* gene under control of a suitable regulatory element, which may be constitutive, tissue specific, developmentally regulated, or inducible and, optionally, a marker gene for selection of transformants;
- ii) selecting transformed plants;
- iii) producing lines that ectopically overexpress the *BNM3* gene, or *BNM3* protein;
- iv) isolating microspores and pollen from the transgenic lines and culturing microspores and pollen to induce embryogenesis.

[0075] Embryogenesis can be induced by any suitable protocol, for example, which is not to be considered limiting, culturing microspore and pollen for about four days at from about 28° to about 35 °C, preferably at about 32°C, then transferring embryogenic cells or embryos to about 25 °C.

[0076] Using the above method, *Brassica napus* cultivars ectopically overexpressing *BNM3* show an increase in the percentage of embryogenic cells or embryos over that observed when microspores or pollen are prepared from wild-type plants that do not ectopically express *BNM3*.

[0077] Examples of regulatory elements that may be used to express *BNM3* for the production of microspore-derived embryos include, but are not limited to, those of the class I low molecular weight heat shock inducible gene, *GMHSP17.3B* (Zarsky *et al.*, 1995, which is incorporated by reference), or microspore/pollen expressed genes such as *NTM19* (Oldenhof *et al.*, 1996, EP 790,311, which are incorporated by reference), *BCP1* (Xu *et al.*, 1995 which is incorporated by reference), *LAT52* (Twell *et al.*, 1989, which is incorporated by reference), *BNM1* (Treacy *et al.* 1997, which is incorporated by reference), and *APG* (Roberts *et al.*, 1993, which is incorporated by reference). Also useful are inducible regulatory elements, for example but not limited to, tetracycline-inducible promoter (Gatz 1997, which is incorporated by reference), steroid inducible promoter (Aoyama and Chua 1997, which is incorporated by reference) and ethanol-inducible promoter (Slater *et al.* 1998, Caddick *et al.* 1998, which are incorporated by reference).

[0078] In a similar fashion, microspore-derived embryos may also be produced in plants by introducing into a plant

of interest a BNM3 protein, (e.g. via biolistics; Klein et al 1987) and selecting for plants that exhibit increased microspore embryogenesis.

[0079] This invention also provides a method for the efficient production of somatic embryos *in vitro*. This method involves:

- i) transforming a plant, for example, *Arabidopsis* using transformation techniques known to one of skill (for example, but not limited to, DeBlock *et al.* 1989, Clough and Bent 1998, Vergunst *et al.* 1998, which are incorporated by reference), or a plant cell may also be transiently transformed using methods known to one of skill (for example, biolistics; Klein et al 1987) with a vector construct containing a *BNM3* gene under control of suitable regulatory element, which may be constitutive, inducible or developmentally regulated, and, optionally, a marker gene for selection of transformants is transformed to several *Arabidopsis*.
- ii) selecting transformed plants, and
- iii) culturing the desired explant from the selected transformed plants, for example, but not limited to, root, leaf or seedlings *in vitro*, in media with or without appropriate growth regulators, for example, but not limited to 2,4-D (e.g. Mordhorst *et al.*, 1998) to produce direct embryogenesis or embryogenic callus; and
- iv) transferring embryos, non-embryogenic callus, or both embryos and non-embryogenic callus to appropriate media for the production of embryos, plantlets, or both embryos or plantlets.

For example, when the results of the above method are compared with the production of somatic embryos *in vitro* using a number of *Arabidopsis* ecotypes, directed embryogenesis or embryogenic callus is initiated at a higher frequency from transgenic lines ectopically over-expressing *BNM3* than in wild-type controls.

[0080] Examples of regulatory elements that may be used to express *BNM3* for the production of somatic embryos include, but are not limited to, those of genes activated by plant growth regulators which are routinely used to induce somatic embryogenesis in tissue culture. Specific examples, which are to be considered non-limiting, cytokinin inducible *IB6* and *CK1* genes (Brandstätter and Kieber, 1996; Kakimoto, 1996, which are incorporated by reference) and the auxin inducible element, *DR5* (Ulmasov *et al.*, 1997, which is incorporated by reference). Also useful are inducible regulatory elements, for example but not limited to, a tetracycline-inducible promoter (Gatz 1997, which is incorporated by reference), a steroid inducible promoter (Aoyama and Chua 1997, which is incorporated by reference), and an ethanol-inducible promoter (Slater *et al.* 1998, Caddick *et al.* 1998, which are incorporated by reference).

[0081] Ectopic initiation of embryo development is one of the key steps in apomixis. As shown in Example 4, ectopic expression of a *BNM3* gene is sufficient to initiate embryo formation in otherwise non-embryo-forming tissue. A *BNM3* gene may therefore be used to initiate adventitious embryony or parthenogenesis of a reduced or unreduced embryo sac cell by expression of the gene in the sporophytic or gametophytic tissues of the developing ovule.

[0082] Adventitious embryony is achieved by expressing *BNM3* in sporophytic ovule tissues such as the nucellus, the inner integuments or other tissues lying adjacent to or in proximity to the developing embryo sac. This method involves:

- i) transforming a desired plant (see above methods) with a vector construct consisting of a *BNM3* gene under control of suitable regulatory element, which may be constitutive, inducible or developmentally regulated, and, optionally, a marker gene for selection of transformants, using methods known within the art;
- ii) selecting transformed plants;
- iii) emasculating the transformed plant;
- iv) pollinating the transformed plants with pollen carrying one or more dominant selectable markers, for example GUS or kanamycin resistance; and
- v) assaying for production of clonal offspring.

When the results of the above method are compared with the pollination of a wild-type *Arabidopsis* plant with pollen carrying the dominant selectable marker, all F1 embryos resulting from this cross inherit the dominant marker while embryos derived from plants ectopically over-expressing the *BNM3* gene or protein are clonally derived via sexual embryo formation and do not inherit the dominant selectable marker.

[0083] Specific examples of gene regulatory elements suitable for directing expression of *BNM3* to obtain adventitious embryony, diplospory or haploid parthenogenesis of embryo sac components include the ovule expressed *SERK* gene (Schmidt *et al.* 1997, which is incorporated by reference), the meiosis expressed *AtDMC1* gene, (Klimyuk and Jones, 1997; WO 98/28431, which are incorporated by reference), the ovule expressed *AGL11* gene (Roundsley *et al.*, 1995, which is incorporated by reference), the nucellus expressed *NUC1* gene (Doan *et al.*, 1996; WO 98/08961, which are incorporated by reference), and the inner integument-expressed genes, *FBP7* (Angenent *et al.* 1995, which is incorporated by reference) and *SC4* (US application 09/059,909, filed April 13, 1998, which is incorporated by reference) genes. Furthermore, inducible systems, for example but not limited to, tetracycline-inducible promoter (Gatz

1997, which is incorporated by reference), steroid inducible promoter (Aoyama and Chua 1997, which is incorporated by reference), ethanol-inducible promoter (Slater et al 1998, Caddick et al. 1998, which are incorporated by reference) may also be used. Parthenogenesis from cells of the embryo sac requires a regulatory element that is active in one or more cells of the female gametophyte or their precursors. Fertilization of the meiotically-derived polar nuclei is desirable when the development of seed is dependent on the presence of endosperm.

Use of *BNM3* Sequences to Control Regeneration Processes

[0084] Plants ectopically over-expressing the *BNM3* genes exhibit increased regenerative capacity and the ability to regenerate whole plants in the absence of added growth regulators (see example 5). *BNM3* gene expression may therefore be used to enhance or induce the regeneration capacity of plant tissues *in vivo* or *in vitro*. The regulatory elements used to express *BNM3* will depend, in part, on the target tissue used for regeneration. Regeneration of plant tissues may be obtained by expressing a *BNM3* gene under the control of a constitutive regulatory element, for example, but not limited to, 35S, or by expressing a *BNM3* gene under the control of tissue specific or developmentally regulated elements, inducible elements derived from either plant or non-plant genes (e.g. Gatz and Lenk, 1998, which is incorporated by reference), or through transient expression methods which do not result in stable integration of the *BNM3* gene or which make direct use of the *BNM3* protein (e.g. microprojectile bombardment of DNA or protein). Chemical induction systems (see Gatz and Lenk, 1998) or regulatory elements of genes that respond to plant growth regulators used to induce regeneration, such as, for example, cytokinin (Brandstatter and Kieber, 1998; Kakimoto, 1996) or auxin (Ulmason et al., 1997), or genes expressed at the wound site of tissue explants (Xu et al., 1993) may be used.

[0085] A further application is the use of a *BNM3* gene as a selectable marker for the recovery of transgenic plants. As an example of this application which is not to be considered limiting in any manner, roots of a seedling, for example, an *Arabidopsis* ecotype C24 seedling, are cocultivated with a single *Agrobacterium tumefaciens* strain (per Vergunst et al, 1998; except that all steps are carried out in the absence of added growth regulators) containing two binary constructs:

- a first binary vector carries a reporter gene fusion, for example, but not limited to, 35S:*GUS*;
- a second binary vector contains a *BNM3* gene under control of suitable regulatory element.

BNM3 gene expression is activated upon integration of the above construct into the arabidopsis genome and transgenic plants are selected on the basis of their ability to regenerate under conditions in which wild-type explants are unable to regenerate, for example, but not limited to, the absence of growth regulators. In many instances the T-DNA carrying the *BNM3* gene and the T-DNA carrying the gene of interest will integrate at unlinked loci. The T-DNA containing the introduced *BNM3* sequence, and it's associated increased regenerative capacity phenotype, may therefore be removed in the progeny plants by simple segregation (Daley et al. 1998). However, as will be apparent to one of skill in the art, other methods such as transient expression, which do not result in stable integration of the *BNM3* gene or which make direct use of the *BNM3* protein, may also be employed.

Use of *BNM3* Sequences to Target Gene Expression to the Embryo

[0086] Since *BNM3* genes are preferentially expressed in developing embryos (see example 3), a further application of this invention is the use of *BNM3* regulatory regions to target expression of at least one heterologous gene of interest to the developing embryo for any purpose, for example, but not limited to, altering embryo and seed traits such as seed viability or size, composition of constituents of the seed, disease resistance, or the production of high value products such as vaccines antibodies, biopharmaceuticals or other specialty chemicals.

Use of *BNM3* Expression as a Marker for Embryogenic Cells

[0087] As shown in Examples 3 and 4, *BNM3* gene expression is detected during the earliest phase of plant embryogenesis and is itself sufficient to activate signal transduction cascades leading to embryo development. *BNM3* gene expression is therefore a specific marker for the entry of a plant cell into the embryogenic pathway.

[0088] *BNM3* expression is associated with embryo-forming cell divisions *in vitro* and *in vivo* and as such can be used to define culture conditions that alter the embryo-forming capacity of a tissue *in vitro*. Cells with embryogenic capacity or cells that undergo only a limited number of embryo-forming divisions are difficult to identify in the absence of structures that morphologically resemble embryos. However, these cells may be identified on the basis of *BNM3* expression. In this application, a vector containing the *BNM3* regulatory region, fused to a reporter gene, for example, but limited to, *GUS* (Jefferson et al., 1987), *Luciferase* (Ow et al., 1987) or *GFP* (Haselhoff and Amos, 1995) is transformed to a plant of interest. Homozygous transgenic lines exhibiting high levels of reporter gene expression in the

embryo are cultured under *in vitro* conditions. Embryogenic cells, as well as culture conditions which facilitate or enhance the formation of embryogenic cells are identified on the basis of reporter gene expression within the cultured tissue.

[0089] A related application is the use of the *BNM3* gene as a marker in apomictic species for the identification of individual cells that are in the process of forming asexually-derived embryos. In this application, cells entering the autonomous embryo pathway are identified by mRNA *in situ* hybridization using a RNA probe derived from a *BNM3* gene sequence, by immunocytochemistry using an antibody directed against a BNM3 protein, by transforming plants with a DNA construct containing a gene fusion between *BNM3* regulatory regions and a reporter gene, or by any similar technique known to those skilled in the art.

Identification of Signal Transduction Components

[0090] Signal transduction components which activate or are activated by *BNM3* gene expression can be elucidated by identifying proteins and DNA sequences that interact with a *BNM3* gene and its protein product. These signal transduction components may be identified using techniques known to a person skilled in the art, including for example, but not limited to:

- yeast one hybrid screens for the isolation of proteins that bind to the *BNM3* regulatory regions to influence *BNM3* gene expression;
- genetic selection in yeast to identify genes that are direct targets of BNM3 binding;
- DNA arrays or proteomics to identify genes which are activated in a BNM3 signal transduction cascade; and
- yeast two hybrid screens to identify proteins that interact with BNM3 to influence expression of downstream target genes.

Techniques for the analysis of the signal transduction components and signalling components are well known (see for example, Meijer *et al.* (1998), Lipshutz *et al.* (1999), and Anderson and Anderson (1998)).

[0091] Plants over-expressing the *BNM3* gene under control of a strong constitutive regulatory element such as, for example, but not limited to, the Cauliflower Mosaic Virus 35S promoter exhibit ectopic embryo formation, enhanced regeneration via organogenesis or a combination thereof (Examples 4 and 5). The ability of *BNM3* ectopic over-expression to induce both embryo formation and enhance regeneration processes can be used to identify mutants altered in their embryo-forming or regenerative capacity. In this application a vector construct consisting of a *BNM3* protein coding region under control of a regulatory element that is sufficient to promote either ectopic embryo formation or enhanced regeneration phenotype is made and introduced into a plant of interest. Homozygous transgenic lines exhibiting a high penetrance of ectopic embryo formation, enhanced regeneration phenotype, or a combination thereof are identified. These lines are mutagenized by any available technique well known to the person skilled in the art, but which may include EMS mutagenesis, fast neutron mutagenesis, transposon mutagenesis or T-DNA mutagenesis. Mutagenized plants are then screened for alterations in the ectopic embryo formation or regeneration phenotype. These alterations include, for example, but not limited to, elimination or enhancement of the ability to promote ectopic asexual embryo formation or to regenerate in the absence of added growth regulators.

Heterologous Protein Expression System

[0092] Genetic control of the signal transduction pathway leading to embryogenesis and organogenesis in non-seed organs of transgenic plants may be activated by ectopic expression of a *BNM3* gene. Expression of a *BNM3* gene in association with a heterologous promoter can be used to produce altered seed components including for example, proteins, oils and other metabolites. Biotransformation of desired organs may also include altering the nutritive value of, for example leaves of forage crops, or it may be used to create alternative uses for crops. The use of promoters that are induced by the signal transduction cascade initiated by expression of *BNM3* can be used to express high-valued recombinant proteins in organs other than seeds. An example of one such promoter is the napin promoter, obtained from the 2S seed storage protein napin. The production of proteins initiated from a *BNM3*-induced cascade, may be achieved within organs exhibiting greater biomass than seeds. Therefore, this technology may be used to create alternatives for plants as crops.

[0093] Accordingly, the present invention further relates to a binary system in which the BNM3 protein binds directly or indirectly to an embryo-expressed regulatory sequence (target sequence) and activate transcription of a chimeric gene construct in any plant cell, tissue or organ. Therefore, BNM3 may be used to directly or indirectly activate transcription of a chimeric gene construct. This approach involves BNM3 interacting either directly with at least one target sequence from an embryo-expressed gene, or indirectly by initiating an embryogenic signal cascade that activates a transcription factor that in turn binds to and activates transcription from at least one target sequence. This binary system

may be used for the expression of proteins in somatic tissues with the properties of expression in seeds.

[0094] In this application transgenic plants containing the *BNM3* gene under control of a constitutive regulatory element, for example, but not limited to the *35S* promoter (*35S:BNM3*) are created to produce a *BNM3* activator line. *BNM3* expression may be demonstrated in a wide range of tissues in the *BNM3* activator lines by RNA gel blot analysis. Stable homozygous activator lines with high levels of *BNM3* expression are identified. Somatic tissues over-expressing *BNM3* may be examined for expression of other embryo-expressed genes, such as arabin (Guerche *et al.*, 1990), cruciferin (Pang *et al.*, 1988) or oleosin, or for morphological properties that are normally characteristic of seeds, such as the presence of lipid or protein bodies.

[0095] Transgenic plants of the same species to that used to generate the *BNM3* activator lines described above are also created which contain an embryo-expressed promoter fused to a gene of interest, to produce a gene of interest line. In order to help describe this embodiment, the gene of interest line expresses a reporter gene, such as *GUS*, and examples, which are not to be considered limiting, of such lines include *Brassica napus* 2S albumin seed storage protein gene, *BngNAP1:GUS* fusion (Baszczynski *et al.*, 1994) or a *SERK:GUS* fusion (Schmidt *et al.*, 1997, a non-seed expressed reporter construct such as *BNM1:GUS* (Treacy *et al.*, 1997) may be used as a negative control). The fidelity of expression of the gene of interest in the specific organs and tissues of these gene of interest lines is demonstrated for each construct. Stable homozygous lines with high levels of expression of the gene of interest expression are created.

[0096] Transgenic lines containing *BNM3* activator lines and gene of interest lines are crossed and the progeny seeds collected. *BNM3* gene expression, and in this example, *GUS* activity, expression of other embryo-expressed genes, as well as the morphological characteristics of transformed tissues, are examined. *BNM3* expression in non-seed tissues typically activates both embryo development and expression of the gene of interest (e.g. *GUS*), however, activation of the expression of the gene of interest in the absence of morphologically discernible embryos may also be observed. Expression of the gene of interest, in the absence of morphologically discernible embryos provides initial evidence for direct interaction of *BNM3* with the target sequence.

[0097] Direct interaction of *BNM3* with a target sequence may also be demonstrated using transient expression of *BNM3* in plant protoplasts, along with the transient co-expression of an embryo-expressed promoter fused to a gene of interest (i.e. a gene of interest construct). *35S:BNM3* DNA and the gene of interest construct are introduced into protoplasts derived from non-seed cells, such as leaf mesophyll cells by electroporation. The expression of the gene of interest is examined after several hours to confirm activation of the target sequence. Direct interaction of *BNM3* with the target sequence may further be demonstrated by co-introducing the target sequence alone as competitor DNA.

[0098] In order to determine if tissues from different plant species may be transactivated by *BNM3*, *35S:BNM3* DNA and a reporter gene (for example, but not limited to *GUS*) construct may be introduced by microprojectile bombardment into somatic tissues of a plant. If *BNM3* interacts directly with a target sequence then expression of the report er gene should coincide with transient expression of *BNM3* in all species and tissues.

[0099] Direct evidence for *BNM3*-target sequence interaction may also be obtained by isolation of *BNM3* protein expressed in bacteria, insect or yeast. *BNM3* is expressed in bacteria, insect, or yeast using commercially available expression systems and isolated to purity. Gel mobility shift assays (Gustavson *et al.*, 1991) are performed using a *BNM3*-target sequence, for example an embryo-expressed target sequence, to demonstrate direct binding of *BNM3* to the *BNM3*-target sequence. Footprint analyses may also be performed to locate the region of *BNM3* binding. Fragments of target sequences that bind *BNM3* may then be subcloned and used as competitors for *BNM3* binding in transient assays described above.

[0100] The present invention will be further illustrated in the following examples. However it is to be understood that these examples are for illustrative purposes only, and should not be used to limit the scope of the present invention in any manner.

Examples

General methods: Microspore Embryo Culture

[0101] *Brassica napus* c.v. Topas was used as the source of all plant material for microspore embryo culture. Donor plants for microspore culture were grown in a growth cabinet at 20°C/15°C (day/night) with a 16 h photoperiod (400 μ E/m/s) provided by VHO cool white fluorescent lamps (165W, Sylvania) and incandescent bulbs (40W, Duro-test). Four weeks after germination the plants were transferred to growth cabinets under the same light conditions, but set at 10°C/5 °C (day/night). Microspores and pollen were isolated and cultured as described in Keller *et al.* (1987), except that after 21 days in culture, cotyledon stage embryos were transferred to a maturation medium consisting of 1/2X NLN salts, 1% sucrose, 0.35 M mannitol and 5 μ M ABA. Uninduced cultures (microspores and pollen continuing gametophytic development) and heat-stressed, non-embryogenic cultures (used for construction of the subtracted probe), were cultured from the same starting material as was used for the initiation of embryogenic cultures. Uninduced

samples were obtained by culturing microspores and pollen for four days at 25 °C. Heat-stressed, non-embryogenic samples were obtained by culturing microspores and pollen for one day 25 °C, followed by three days 32°C.

[0102] Samples of microspore and pollen cultured for less than 10 days were collected by centrifugation. Older samples containing globular, heart, torpedo and cotyledon stage microspore-derived embryos were collected by filtration through nylon meshes of various pore sizes as described in Ouellet *et al.* (1992). All other plant tissues were collected from greenhouse grown material. Seed material was obtained by hand pollinating flowers on the day of anthesis and collecting developing seeds on various days after pollination (DAP).

Nucleic Acid Isolation and Analysis

[0103] Total RNA was isolated using either a cesium chloride/guanidinium isothiocyanate procedure (Ouellet, 1992) or TRIzol reagent (Gibco-BRL). RNA gel blot analysis was carried out by separation of 5 to 20 µg of total RNA per lane through 1.5% agarose gels containing 0.62 M formaldehyde, essentially as in Sambrook *et al.* (1989), followed by capillary transfer to Hybond-N nylon membranes (Amersham). Poly(A)⁺ RNA was isolated from total RNA by oligo (dT)-cellulose chromatography (Sambrook, 1989).

[0104] Genomic DNA was isolated from leaf tissue as described in Fobert *et al.* (1991) and digested with the specified restriction enzymes using standard procedures (Sambrook, 1989). DNA gel blot analysis was carried out by electrophoresis of 10 µg DNA through 0.8% agarose gels followed by capillary transfer to Hybond-N membranes.

[0105] The partial 1.2 kb *BNM3A* cDNA insert was used as a probe for DNA and RNA gel blots. Hybridization to gel blots was carried out at 65 °C according to the Hybond-N protocol. The final wash conditions were 0.1X SSC, 65°.

Subtractive Probe Construction and cDNA Library Screening

[0106] Poly (A) mRNA was isolated from late uninucleate microspores and early binucleate pollen that had been cultured for four days at 32 °C in order to induce embryogenesis (embryogenic sample) and used to synthesize first strand cDNA (Riboclone cDNA kit; Promega). The cDNA was then hybridized to a five-fold excess (by weight) of poly (A)⁺ RNA from late uninucleate microspores and early binucleate pollen that had been cultured for one day at 25 °C, followed by three days at 32 °C to inactivate embryogenesis (non-embryogenic sample; Pechan *et al.* 1991). The subtractive hybridization was performed essentially as described in Sambrook *et al.* (1989). The single-stranded cDNA recovered after subtraction was labelled with [α -³²P] dCTP using a random primers kit (BRL) and used as the subtracted probe for screening a Lambda phage cDNA library constructed from the same embryogenic sample described above (Boutillier, 1994). Triplicate nylon filter lifts (Hybond-N) from approximately 1.5 x 10⁶ plaque-forming units of the library were screened with the subtracted probe, with a random primers-labelled first strand non-embryogenic probe and with a random primers-labelled napin seed storage protein cDNA probe (pN2; (Crouch, 1983). Napin mRNAs are prevalent in the embryogenic microspore library (Boutillier, 1994) and therefore plaques hybridizing to the napin probe were removed from the subsequent screening steps. Plaques hybridizing to the subtracted probe, but not to the non-embryogenic or napin probes, were selected and subjected to two subsequent rounds of differential screening using both the subtracted and non-embryogenic cDNA probes. DNA from selected Lambda clones was isolated (Sambrook, 1989), partially digested with *Eco* RI and *Xba* I and subcloned into pGEM-4Z (Promega).

[0107] Seven cDNAs comprising 6 unique genes, one of which comprised a truncated *BNM3A* cDNA, were identified. Two distinct, full length *BNM3* cDNA clones (*BNM3A* and *BNM3B*) were subsequently obtained by stringent screening of circa 2.5 x 10⁶ plaque-forming units of a cDNA library (UniZAPIII cDNA synthesis kit, Stratagene) constructed with mRNA from 10 day old globular to heart-stage microspore-derived embryos of *B. napus* c.v. Topas. The *BNM3* cDNA inserts were rescued by *in vivo* excision into Bluescript SK(-) (Stratagene).

Isolation of Genomic DNA sequences

[0108] The Universal Genome Walker Kit (Clontech) was used to isolate genomic DNA fragments lying upstream of the *BNM3* ATG start codon. Pools of uncloned, adaptor-ligated *Brassica napus* cv Topas genomic DNA fragments were constructed and used to isolate *BNM3* genomic sequences by nested PCR. The primary PCR made use of the outer adaptor primer (AP1) supplied by the manufacturer and a *BNM3* specific primer with the sequence:

5'-GAGGCAGCGGTGCGATCGTAACAGTACTCT-3' (SEQ ID NO:6).

The nested PCR made use of the nested adaptor primer (AP2) supplied by the manufacturer and a *BNM3* specific primer with the sequence:

5'-CATAAGGAGAGAGAGAAAAGCCTAACCACT-3' (SEQ ID NO:7).

The primary PCR mixture was then diluted 1:50 and used as template for nested PCR. Both the primary and nested PCR products were performed as recommended by the manufacturer. The nested PCR products were cloned into the pGEMT-Easy vector (Promega) and sequenced. PCR products corresponding to the 5' untranslated genomic regions of both *BNM3A* and *BNM3B* cDNAs were identified.

[0109] The genomic DNA sequence spanning the *BNM3A* ATG translational start and TAG translational stop codons was isolated by PCR from *B. napus* cv Topas genomic DNA using *Pfu* polymerase (Stratagene) and the following primer combination:

5'-ACCAAGAAGCTCGTTAGATC-3' (SEQ ID NO:8);

and

5'-AACGCATATAACTAAAGATC-3' (SEQ ID NO:9).

The primers were used under standard PCR conditions. The PCR products were cloned into the pGEMT-Easy vector and sequenced.

Plasmid Construction for Plant Transformation

[0110] The construction of a plasmid vectors containing the *BNM3* cDNAs under control of either a *POLYUBIQUITIN* or Cauliflower Mosaic virus 35S promoter are described below. The plasmid pRAP2TUBI contains a modified *Helianthus annuus* *POLYUBIQUITIN* promoter (Binet *et al.*, 1991) in the plasmid pRAP2T. The plasmid pRAP2T consists of the pUCAP plasmid (van Engelen *et al.*, 1995) and a nopaline synthase (*nos*) terminator inserted into the *Sac* I and *Eco* RI restriction sites. A PCR fragment of the *POLYUBIQUITIN Ubb1* promoter comprising the 5' end of the promoter to 7 bp from the 3' end of the first exon was amplified from the vector using an M13 reverse primer and the UBIQ-3' primer:

5'-CCATGGATCCAGAGACGAAGCGAAAC-3' (SEQ ID NO:10)

which includes introduced *Nco* I and *Bam* HI restriction sites. The *POLYUBIQUITIN* promoter fragment was digested with *Pst* I and *Bam* HI, gel purified and ligated into the *Pst* I and *Bam* HI sites of pRAP2T, creating the vector pRAP2TUBIHa. The full-length *BNM3B* cDNA was digested with *Eco* RI and *Xho* I restriction enzymes, blunted with Klenow enzyme, gel purified and ligated into the *Sma* I site of pRAP2TUBI making the plasmid pKB1S. An *Asc* I/*Pac* I DNA restriction fragment containing the modified *POLYUBIQUITIN* promoter, the *BNM3B* cDNA and the *nos* terminator was gel purified, and ligated to the *Asc* I/*Pac* I digested binary vector pBINPLUS (van Engelen *et al.*, 1995), creating the plasmid pKBBIN1S.

[0111] The construction of a vector containing the *BNM3A* cDNA under control of a double enhanced 35S promoter and AMV translational enhancer was as follows. A *Hind* III/*Xba* I DNA restriction fragment containing the double 35S promoter and the AMV translational enhancer from plasmid pB1525 (Datta *et al.*, 1993) was ligated to *Hind* III/*Xba* I digested pRAP2T, creating the plasmid pRAP2T35S. An *Nco* I site was introduced into the *BNM3A* cDNA clone by site directed mutagenesis. The sequence of the BNM3ANCO1 primer used for mutagenesis is:

5'-ACTCCATGGATAATAACTGGTTAGGC-3' (SEQ ID NO:11).

A second primer, BNM3AHINDIII:

5' - AAATTCTCAAGCTTTGGTCCATCTTG-3' (SEQ ID NO12)

was used together with the BNM3ANCO1 primer to amplify a 305 bp fragment of the *BNM3A* cDNA. This PCR fragment was digested with *Nco* I and *Hind* III and ligated to *Nco* I/*Kpn* I cut pRAP2T35S and a *Hind* III/*Kpn* I fragment containing the region of the *BNM3A* cDNA downstream of the *Hind* III site, creating the vector p35S:BNM3. p35S:BNM3 was

digested *Asc*I and *Pac*I restriction enzymes and the fragment containing the double 35S promoter, the AMV translational enhancer, the BNM3A cDNA and the nos terminator was gel purified and ligated to the *Asc*I/*Pac*I digested binary vector pBINPLUS, creating the plasmid p35S:BNM3BIN.

[0112] Both the pKBBIN1S and p35S:BNM3BIN plasmids were transferred to *Agrobacterium tumefaciens* C58C1 strain carrying the disarmed Ti plasmid pMP90 and used in transformation experiments.

Plant Transformation

[0113] *Arabidopsis thaliana* ecotype C24 was used as the recipient in transformation experiments. Plants were transformed using either the floral dip method described in Clough and Bent (1998) or the root transformation method described in Vergunst *et al.* (1998).

[0114] Transgenic *Brassica napus* c.v. "Topas" plants were produced by *Agrobacterium tumefaciens*-mediated transformation of microspore-derived embryos. Microspore-derived embryos were cultured for 5 weeks at a density of approximately 1000 embryos per ml. Overnight cultures of *Agrobacterium* were diluted 100 times in B5 medium containing 9% sucrose. Embryos were co-cultivated with the diluted bacteria for 48 hours at 24°C in darkness, with slow shaking. The embryos were then transferred to NLN13 medium supplemented with 350 mg/L cefotaxim and 200 mg/L vancomycin for at least two weeks in darkness at 25 °C.

[0115] Embryos were germinated in weak light at 25°C for about 2 weeks on solid B5 medium supplemented with 2% sucrose, cefotaxim (200 mg/L) and vancomycin (100 mg/L). Well developed hypocotyls from germinated embryos were isolated and transferred to fresh germination medium supplemented with 100 mg/L kanamycin. After two weeks on this medium, explants were subcultured to a similar medium supplemented with kanamycin (25 mg/L). Green, putative transgenic, secondary embryos become visible after one month of selection.

Microscopy

[0116] All plant material was fixed overnight at 4 °C in 0.1 M phosphate buffer pH 7.0 containing 4% paraformaldehyde. Samples were washed in 0.1 M phosphate buffer and then dehydrated in a graded ethanol series to 100% ethanol. Samples for scanning electron microscopy were critical point dried in liquid CO₂ (Balzers CPD020), and mounted on SEM stubs using conductive carbon glue. Samples were coated with 30 nm palladium/gold using a Polaron E5100 sputter coater. Samples were observed in a JEOL JSM 5200 scanning electron microscope with an acceleration voltage of 15 kV. Digital images were obtained using Orion Framegrabber. Samples for light microscopy were embedded in Technovit 7100 (Kulzer). Sections were stained for 10 seconds in 1% Toluidine blue in 1% sodiumtetraborate, rinsed with water and mounted in Euparal. Digital images were recorded using a Sony 3 CCD camera.

Regeneration Experiments

[0117] Wild-type and transgenic arabidopsis seeds were surface sterilized, plated on 1/2 MS media containing 20% sucrose (1/2MS-20) and grown at 21°C with the plates inclined at a 60° angle. Eight wild-type seedlings and eight seedlings from each of seven independent transgenic lines were harvested 10 days after germination and separated into root, hypocotyl and leaf explants. This material was then divided into two batches. Half of the explants were continuously cultured on B5 media containing 20% glucose (B5-20). Explants were transferred to fresh B5-20 media every two weeks. The remaining explants were cultured on B5-20 containing plant growth regulators in order to induce shoot regeneration (Vergunst *et al.* 1998). These explants were first placed on callus inducing media (CLM; high auxin to cytokinin ratio) for two days and then transferred to shoot inducing media (SIM; high cytokinin to auxin ratio) for the remainder of the culture period. Explants were transferred to fresh SIM media every two weeks.

Example 1: Isolation and Characterization of the BNM3 Genes from *Brassica napus*

[0118] A subtractive screening approach was used to isolate genes preferentially expressed during the induction of *Brassica napus* c.v. Topas microspore embryogenesis (Figure 1). Two types of microspore cultures were used in the construction of a subtracted probe: embryogenic and non-embryogenic. Embryogenic cultures were obtained by subjecting late uninucleate microspores and early binucleate pollen to a 4 day, 32 °C heat stress treatment. The non-embryogenic sample was obtained by culturing the same starting population of late uninucleate microspores and early binucleate pollen for 1 day at 25 °C followed by 3 days at 32 °C (Pechan *et al.*, 1991). Poly(A) mRNA was isolated from the embryogenic sample and used to synthesize first strand cDNA. The cDNA was then hybridized to an excess of poly(A)⁺ RNA isolated from a non-embryogenic microspore/pollen sample. The non-hybridizing, single stranded cDNA, enriched for sequences present in the embryogenic sample, but absent or present at a much lower level in the non-embryogenic sample, was recovered, radioactively labelled and used as a subtracted probe for screening a cDNA

library derived from the embryogenic sample described above. Plaques hybridizing to the subtracted probe, but not to a probe derived from the non-embryogenic sample, were selected and subjected to two subsequent rounds of differential screening. Seven independent cDNA clones, comprising six unique DNA sequences were found to be differentially expressed between the embryogenic and non-embryogenic samples. One of these clones, 42A1, later renamed *BNM3A* (for *Brassica napus* microspore embryo), was further characterized.

Example 2: The BNM3 genes encode new members of the AP2 domain class of transcriptional activators

[0119] A single *BNM3* cDNA clone, *BNM3A*, was isolated after screening an embryogenic microspore cDNA library with a subtracted probe enriched for genes expressed in embryogenic microspores and pollen. The discrepancy between the size of the cDNA clone (1.2 kb) and the size of the transcript detected on RNA gel blots (2.2 kb) indicated that this clone did not represent a full-length cDNA. Two longer cDNA clones, corresponding to the full length cDNA of the clone originally isolated, *BNM3A* (SEQ ID NO. 1), and a new clone, *BNM3B* (SEQ ID NO. 3), were isolated from a 10 day old *Brassica napus* microspore embryo cDNA library. The alignment of the DNA sequence of these clones is shown in Figure 2. The two *BNM3* cDNA clones are 2011 and 1992 nt in length, and are 97% similar at the nucleotide level, differing only slightly in the length and sequence of their 5' and 3' untranslated regions. Both cDNAs potentially encode 579 amino acid polypeptides (predicted molecular mass of 63.9 kDa, pI of 5.7) that are 97% similar at the amino acid level (Figure 3).

[0120] The genomic complexity of the *BNM3* genes was determined by hybridization of the *BNM3* cDNAs to gel blots containing *B. napus* genomic DNA (Figure 4). The *BNM3* cDNAs hybridize to two DNA fragments under high stringency conditions. The two hybridizing fragments represent the two *BNM3* genes, *BNM3A* and *BNM3B*. *B. napus* is an amphidiploid species derived from the hybridization of the diploid *B. rapa* and *B. oleracea* genomes, thus the two *BNM3* sequences are likely derived from a single copy locus in each of the parental diploid progenitors.

[0121] Search of the sequence databases indicated that the *BNM3* translation products contain two copies of an AP2 domain (Figure 3). The AP2 domain was first identified in APETALA2 (AP2), an arabinopsis protein that regulates meristem identity, floral organ specification, seedcoat development and floral homeotic gene expression (Jofuku *et al.*, 1994; WO 98/07842), and has since been identified in a wide range of proteins with diverse functions. These functions range from the activation of genes involved in stress (Zhou, 1997; Stockinger, 1997) and ethylene response (Ohme-Takagi, 1995) to the regulation of leaf, floral and ovule development (Moose, 1996; Jofuku, 1994; Elliot, 1996; Klucher, 1996). The AP2 domain is a 56-68 amino acid repeated motif containing at least two conserved regions: a highly basic YRG element, containing a conserved YRG amino acid motif and the RAYD element. The RAYD element contains a conserved central core of 18 amino acids that is predicted to form an amphipathic α -helix, a structure that is thought to mediate protein-protein interactions. The ability of a number of AP2 domain containing proteins to bind DNA, coupled with the presence of putative nuclear localization signals and acidic regions that may function as transcriptional activators suggests these proteins function as transcription factors.

[0122] Two phylogenetically distinct classes of AP2 domain proteins, consisting of either one AP2 domain (EREBP-like) or two AP2 domains connected by a linker region (AP2-like), have been identified (Zhou, 1997). *BNM3* belongs to the latter class. Search of the databases with the region corresponding to the two AP2 domains and linker region of *BNM3* reveals that *BNM3* is most similar to the arabinopsis AINTEGUMENTA (ANT; Elliot, 1996; Klucher, 1996) and the *Zea mays* ZMMHCF1 AP2 domain containing protein (ZM; Daniell, 1996) Figure 5 shows an alignment of the two AP2 domains of *BNM3* with those of other proteins that contain two AP2 domains. *BNM3* shares 85% amino acid sequence similarity with ANT and 86% with ZMMHCF1 in this region, but only 66% amino acid similarity with AP2 and GLOSSY15 in this region. A 10 amino acid insertion in the first AP2 domain of the and *BNM3* proteins further distinguishes these three proteins from other AP2 domain containing proteins (Elliot, 1996). The *BNM3*, AINTEGUMENTA and ZMMHCF1 proteins also share a small hydrophobic amino acid motif, LG/SFSL, in their amino terminal regions, but otherwise show no significant similarity in their DNA or amino acid sequences outside of the AP2 domains and linker. These results indicate that the *BNM3* sequences encode unique members of the AP2 domain family of proteins.

[0123] A pairwise alignment of *BNM3B* cDNA and amino acid, sequences with ANT or ZMMHCF1 sequences indicated that for the *BNM3B* nucleotide sequence:

- there is a 56% identity with ANT cDNA (over the 1905 nucleotides of ANT) and a 58% identity with ZMMHCF1 cDNA (over the 1773 nucleotide sequence of ZM);

and for the *BNM3B* amino acid sequence:

- there is a 41% identity of the *BNM3B* protein with ANT protein (over the 555 amino acid sequence of ANT), and a 46% identity with ZMMHCF1 protein (over 485 amino acid sequence of ZM).

Example 3: The BNM3 genes are preferentially expressed in developing embryos

[0124] RNA gel blot analysis (Figure 6) was used to determine the pattern of *BNM3* gene expression during microspore-derived embryo development, seed development, and in non-seed tissues. Both analyses indicate that the *BNM3* genes are preferentially expressed in developing embryos.

[0125] RNA gel blot analysis indicates that *BNM3* mRNAs are detected in microspore cultures induced to undergo embryogenesis, as well as in the subsequent globular, heart, torpedo and cotyledon stages of microspore-derived embryo development (Figure 6A). *BNM3* mRNAs are not detected in non-embryogenic microspore cultures, in freshly isolated microspores and pollen, or in microspores and pollen continuing gametophytic development in culture (Figure 6A). RNA gel blot analysis of developing seeds shows that *BNM3* expression is first detected 14 days after pollination (14 DAP), corresponding to the heart stage of embryo development. *BNM3* expression increases during the early (21 DAP) and mid-cotyledon (28 DAP) stages of embryo development and remains constant thereafter (Figure 6B). *BNM3* transcripts were not detected in any of the non-seed tissues tested, reflecting the low level or absence of transcripts in these tissues.

Example 4: Expression of BNM3 in Vegetative Tissues Promotes Asexual Embryo Formation

[0126] In order to determine the function the *Brassica napus* *BNM3* proteins, the *BNM3* cDNAs were placed under the control of two separate constitutive promoter constructs, a modified sunflower *POLYUBIQUITIN* promoter construct (hereafter referred to as *UBI:BNM3*) and a double enhanced *35S* promoter construct containing an AMV translational enhancer (hereafter referred to as *35S:BNM3*), and introduced into *Arabidopsis*. Analysis of the phenotype of the transformants indicates that ectopic over expression of the *BNM3* cDNAs promotes the formation of somatic embryos on vegetative structures such as cotyledons, petioles, leaf blades and the shoot apical meristem (Figure 7). The frequency of transformants producing ectopic embryos, as well as the penetrance of the ectopic embryo phenotype, was greater when the *BNM3* gene was expressed under control of the stronger double enhanced *35S* promoter-AMV translational enhancer, as compared to the *POLYUBIQUITIN* promoter. Thus a high threshold level of protein product is required to increase the frequency and penetrance of the ectopic embryo phenotype.

[0127] *BNM3*-derived ectopic embryos contain all of the organ systems and tissue layers found in the developing zygotic embryo. *BNM3*-derived ectopic embryos are bipolar (Figures 7D and E) and consist of an axis, comprising of the hypocotyl and radicle regions, shoot and root meristems, and cotyledons (Figure 7E). In addition, each organ system contains the characteristic radial arrangement of three specialized tissue layers (epidermis, ground parenchyma and provascular tissue) found in zygotic embryos (Figure 7E). Continued expression of the *BNM3* gene within the developing ectopic embryo leads to a reiteration of the embryo-forming process, with the result that new embryos are continuously formed on the surface of pre-existing embryos (Figure 7D and E). These results provide conclusive evidence that expression of a single gene, *BNM3*, is sufficient to initiate a signal transduction cascade leading to the formation of fully differentiated asexually-derived embryos.

Example 5: Expression of BNM3 Increases the Regeneration Capacity of Plant Tissues

[0128] We examined the effect of *BNM3* gene expression on the ability of *Arabidopsis* plants to regenerate shoots *in vitro* in the presence or absence of added growth regulators. Leaf, root and hypocotyl explants from 10 day old seedlings of wild-type *Arabidopsis* and transgenic *Arabidopsis* lines expressing *BNM3* under control of the *POLYUBIQUITIN* promoter were placed on media containing growth regulators to induce first callus formation and then shoot organogenesis. Root explants from transgenic lines show at least a 5-fold increase in shoot regeneration in the presence of hormones as compared to wild-type root explants. (Figure 8A). These shoots also developed faster in the transgenic explants as compared to the wild-type. Wild-type leaf and hypocotyl explants responded by producing callus on the cut end of the petiole (Figure 8B). In contrast, explants from transgenic lines immediately produced new shoots (Figure 8B) or roots from the cut end of the petiole. Transgenic explants that initially produced roots eventually also produced shoots.

[0129] Transgenic explants were also able to regenerate in absence of added growth regulators. Wild-type leaf and hypocotyl explants placed on medium lacking growth regulators occasionally produced callus or roots at the cut end of the leaf petiole, however shoots did not regenerate from these structures (Figure 8C,D). Wild-type roots greened and formed thickened nodule-like structures at the junction with lateral roots, but did not develop further. In contrast, transgenic explants placed on media lacking growth regulators regenerated shoots either from the cut end of the leaf and hypocotyl explants or from the nodule-like structures of root explants (Figure 8C,D).

[0130] The present invention has been described with regard to preferred embodiments. However, it will be obvious to persons skilled in the art that a number of variations and modifications can be made without departing from the scope of the invention as described herein.

References

- [0131] Anderson, N.L. and Anderson, N.G. (1998). Proteome and proteomics: new technologies, new concepts and new words. *Electrophoresis* 19, 1853-1861.
- 5 [0132] Angenent, G.C., Franken, J., Busscher, M., van Dijken, A., van Went, J.L., Dons, H.J.M. and van Tunen, A. J. (1995). A novel class of MAD5 box genes involved in ovule development in *Petunia*. *Plant Cell* 7, 1569-1582.
- [0133] Aoyama, T. and Chua, N.H. (1997). A glucocorticoid-mediated transcriptional induction system in transgenic plants. *Plant J.* 2, 397-404.
- 10 [0134] Baszczynski, C.L., and Fallis, L. (1990). Isolation and nucleotide sequence of a genomic clone encoding a new *Brassica napus* napin gene. *Plant Mol. Biol.* 14, 633-635.
- [0135] Binet, M.N., Lepetit, M., Weil, J.H. and Tessier, L.H. (1991). Analysis of a sunflower polyubiquitin promoter by transient expression. *Plant Sci.* 79, 87-94.
- 15 [0136] Boutilier, K.A., Gines, M.J., Demoor, J.M., Huang, B., Baszczynski, C.L., Iyer, V.N. and Miki, B.L. (1994). Expression of the BrnNAP subfamily of napin genes coincides with the induction of *Brassica* microspore embryogenesis. *Plant Mol. Biol.* 26, 1711-1723.
- [0137] Brandstatter, I. and Kieber, J.J. (1998). Two genes with similarity to bacterial response regulators are rapidly and specifically induced by cytokinin in *Arabidopsis*. *Plant Cell* 10, 1009-1019.
- 20 [0138] Caddick, M.X., Greenland, A.J., Jepson, I., Krause, K.P., Qu, N., Riddell, K.V., Salter, M.G., Schuch, W., Sonnewald, U., and Tornsett, A.B. (1998). An ethanol inducible gene switch for plants used to manipulate carbon metabolism. *Nature Biotech.* 16, 177-180.
- [0139] Chaudury, A.M., Letham, D.S., Craig, S. and Dennis, E.S. (1993). *amp-l-a* mutant with higher cytokinin levels and altered embryonic pattern, faster vegetative growth, constitutive photomorphogenesis and precocious flowering. *Plant J.* 4, 907-916.
- 25 [0140] Clough, S.J. and Bent, A.F. (1998). Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J.* 16, 735-743.
- [0141] Custers, J.B.M., Oldenhof, M.T., Schrauwen, J.A.M., Cordewener, J.H.G., Wullems, G.J. and van Lookeren Campagne, M.M. (1997) Analysis of microspore-specific promoters in tobacco. *Plant Mol. Biol.* 35, 689-699.
- [0142] Crouch, M.L., Tenberge, K.M., Simon, A.E. and Ferl, R. (1983). cDNA clones for *Brassica napus* seed storage proteins: evidence from nucleotide sequence analysis that both subunits of napin are cleaved from a precursor polypeptide. *J. Mol. Appl. Genet.* 2:273-283.
- 30 [0143] Daley, M., Knauf, V.C., Summerfelt, K.R. and Turner, J.C. (1998). Co-transformation with one *Agrobacterium tumefaciens* strain containing two binary plasmids as a method for producing marker free transgenic plants. *Plant Cell Rep.* 17, 489-496.
- [0144] Daniell, T.J., Fordham-Skelton, A.P., Vergani, P. and Edwards, R. (1996). Isolation of a maize cDNA (accession no. Z47554) (PGR 96-013) encoding APETALA-2-like binding domains by complementation cloning of an L-isospartyl methyltransferase-deficient mutant of *Escherichia coli*. *Plant Phys.* 110, 1435.
- 35 [0145] Datla, R.S.S., Bekkaoui, F., Hammerlindl, J.K., Pilate, G., Dunstan, D.I. and Crosby, W.L. (1993). Improved high-level constitutive foreign gene expression in plants using an AMV RNA4 untranslated leader sequence. *Plant Sci.* 94, 139-149.
- 40 [0146] DeBlock, M. DeBrower, D. and Tenning, P. (1989). Transformation of *Brassica napus* and *Brassica oleracea* using *Agrobacterium tumefaciens* and the expression of the bar and neo genes in the transgenic plants. *Plant Physiol.* 91: 694-701.
- [0147] Dellaert, L.M.W. (1981) Comparison of X-ray and fast neutron-induced mutant spectra. Experiments in *Arabidopsis thaliana* (L.) Heynh. *Arabidopsis Inf. Ser.*, 18, 16-36.
- 45 [0148] Doan, D. N. P., Linnestad, C., and Olsen, O.-A. (1996). Isolation of molecular markers from the barley endosperm coenocyte and the surrounding nucellus cell layers. *Plant Mol. Biol.* 31: 877-886.
- [0149] Elliot, R.C., Betzner, A.S., Huttner, E., Oakes, M.P., Tucker, W.Q.J., Gerentes, D., Perez, P. and Smyth, D.R. (1996). *AINTEGUMENTA*, an APETALA2-like gene of *Arabidopsis* with pleiotropic roles in ovule development and floral organ growth. *Plant Cell* 8, 155-168.
- 50 [0150] Federoff, N., Furtak, D., and Nelson O. (1984). Cloning of the bronze locus in maize by a simple and general procedure using the transposable controlling element *Ac*. *Proc. Natl. Acad. Sci. USA* 81, 3825-3829.
- [0151] Feldman, K.A., Marks, M.D., Christianson, M.L., and Quatrano, R.S. (1989). A dwarf mutant *Arabidopsis* generated by T-DNA insertion mutagenesis. *Science* 243, 1351-1354.
- 55 [0152] Fobert, P.R., Miki, B.L., and Iyer, V.N. (1991). Detection of gene regulatory signals in plants revealed by T-DNA-mediated fusions. *Plant Mol. Biol.* 17, 837-851.
- [0153] Gatz, C. (1997). Chemical control of gene expression. *Ann. Rev. Plant Physiol. Plant Mol. Biol.* 48, 89-108.
- [0154] Gatz, C. and Lenk, I.R.P. (1998). Promoters that respond to chemical inducers. *Trends Plant Sci.* 3, 352-358.
- [0155] Guerche, P., Tire, C., Grossi De Sa, F., De Clercq, A., Van Montagu, M. and Krebbers, E. (1990). Differential

expression of the Arabidopsis 2S albumin genes and the effect of increasing gene family size. *Plant Cell* 2, 469-478.

[0156] Gustavsson, H.O., Ellerström, M., Stulberg, K., Ezcurra, I., Komán, A., Hoglund, A., Rask, L. and Josefsson, L.-G. (1991). Distinct sequence elements in a napin promoter interact in vitro with DNA-binding proteins from *Brassica napus*. *Physiol. Plant* 82, 205-212.

[0157] Haseloff, J. and Amos, B. (1995). GFP in plants. *Trends Genet.* 11, 328-329.

[0158] Jefferson, R.A. and Bicknell, R. (1996). The potential impacts of apomixis: a molecular genetics approach. In *The Impact of Plant Molecular Genetics*, B.W.S. Sobral, ed (Boston: Birkhäuser), pp. 87-101.

[0159] Jefferson, R.A., Kavanagh, T.A. and Bevan, M.W. (1987). GUS fusions: β -glucuronidase as a sensitive and versatile gene fusion marker in higher plants. *EMBO J.* 6, 3901-3907.

[0160] Jofuku, K.D., den Boer, B.G.W., van Montagu, M. and Okumuro, J.K. (1994). Control of Arabidopsis flower and seed development by the homeotic gene APETALA2. *Plant Cell* 6, 1211-1225.

[0161] Kakimoto, T. (1996). CKI7, a histidine kinase homolog implicated in cytokinin signal transduction. *Science* 274, 982-985.

[0162] Keller, W.A., Fan, Z., Pechan, P., Long, N., Grainger, J. (1987). An efficient method for culture of isolated microspores of *Brassica napus*. *Proceedings of the 7th International Rapeseed Congress*. Poznan, Poland. Vol. 1, 152-157.

Klein, T.M., Wolf, E.D., Wu, R. and Sanford, J.C. (1987). High-velocity microprojectiles for delivering nucleic acids into living cells. *Nature* 327, 70-73

[0163] Klimyuk, V.I. and Jones, J.D.G. (1997). AtDMC1, the Arabidopsis homologue of the yeast DMC1 gene: characterization, transposon-induced allelic variation and meiosis-associated expression. *Plant J.* 11, 1-14.

[0164] Klucher, K.M., Chow, H., Reiser, L. and Fischer, R.L. (1996). The AINTEGUMENTA gene of Arabidopsis required for ovule and female gametophyte development is related to the floral homeotic gene APETALA2. *Plant Cell* 8, 137-153.

[0165] Koltunow, A.M., Bicknell, R.A. and Chaudhury, A.M. (1995). Apomixis: molecular strategies for the generation of genetically identical seeds without fertilization. *Plant Physiol.* 108, 1345-1352.

[0166] Koltunow, A.M. (1993). Apomixis: embryo sacs and embryos formed without meiosis or fertilization in ovules. *Plant Cell* 5, 1425-1437.

[0167] Korneef, M., Hanhart, C.J. and Thiel, F. (1989). A genetic and phenotypic description of *eceriferum* (cer) mutants in Arabidopsis thaliana. *J. Hered.* 80, 118-122.

[0168] Lightner, J., and Casper, T. (1988) Seed Mutagenesis of Arabidopsis. In *Arabidopsis Protocols*, J.M. Martinez-Zapater and J. Salinas eds (Totowa, USA: Humana Press).

[0169] Lipschultz, R.J., Fodor, S.P.A., Gingeras, T.R. and Lockhart, D.J. (1999). High density synthetic oligonucleotide arrays. *Nature Genetics* 21, 20-24.

[0170] Lotan, T., Ohto, M., Matsudaira Yee, K., West, M.A.L., Lo, R., Kwong, R.W., Yamagishi, K., Fischer, R.L., Goldberg, R.B. and Harada, J.J. (1998). Arabidopsis LEAFY COTYLEDON1 is sufficient to induce embryo development in vegetative cells. *Cell* 93, 1195-1205.

[0171] Meijer, A.H., Ouwkerk, P.B.F. and Hoge, J.H.C. (1998). Vectors for transcription factor cloning and target site identification by means of genetic selection in yeast. *Yeast* 14, 1407-1415.

[0172] Moose, S.P. and Sisco, P.H. (1996). Glossy15, an APETALA2-like gene from maize that regulates leaf epidermal cell identity. *Genes Dev.* 10, 3018-3027.

[0173] Mordhorst, A.P., Toonen, M.A.J. and de Vries, S.C. (1997). Plant embryogenesis. *Crit. Rev. Plant Sci.* 16, 535-576.

[0174] Mordhorst, A.P., Voerman, K.J., Hartog, M.V., Meijer, E.A., van Went, J., Koornneef, M., and de Vries, S.C. (1996). Somatic embryogenesis in Arabidopsis thaliana is facilitated by mutations in genes repressing meristematic cell divisions. *Genetics* 149, 549-563.

[0175] Ogas, J., Cheng, J.-C., Sung, R.Z. and Somerville, C. (1997). Cellular differentiation regulated by gibberellin in the Arabidopsis thaliana pickled mutant. *Science* 277, 91-94.

[0176] Ohme-Takagi, M. and Shinshi, H. (1995). Ethylene-inducible DNA binding proteins that interact with an ethylene responsive element. *Plant Cell* 7, 173-182.

[0177] Oldenhof, M.T., de Groot, P.F.M., Visser, J.H., Schrauwen, J.A.M. and Wullems, G.J. (1996). Isolation and characterization of a microspore-specific gene from tobacco. *Plant Mol. Biol.* 31, 213-225.

[0178] Ouellet, T., Rutledge, R.G. and Miki, B.L. (1992). Members of the acetoxyhydroxyacid synthase multigene family of *Brassica napus* have divergent patterns of expression. *Plant J.* 2, 321-330.

[0179] Ow, D.W., Jacobs, J.D. and Howell, S.H. (1987). Functional regions of the cauliflower mosaic virus 35S RNA promoter determined by use of the firefly luciferase gene as a reporter for promoter activity. *Proc. Natl. Acad. Sci. USA* 84, 4870-4874.

- [0180] Ozias-Akins, P., Lubbers, E.L., Hanna, W.W. and McNay, J.W. (1993). Transmission of the apomictic mode of reproduction in *Pennisetum*: co-inheritance of the trait and molecular markers. *Theor. Appl. Genet.* 85, 632-638.
- [0181] Pang, P., Pruitt, R. and Meyerowitz, E. (1988). Molecular cloning, genomic organisation, expression and evolution of the 12S storage protein genes of *Arabidopsis thaliana*. *Plant Mol. Biol.* 11, 805-820.
- [0182] Peacock, W.J., Ming, L., Craig, S., Dennis, E., Chaudury, A.M. (1995). A mutagenesis programme for apomixis genes in *Arabidopsis*. In *Proceedings Symposium on Induced Mutations and Molecular Techniques for Crop Improvement*. (Vienna: International Atomic Energy Agency), pp. 117-125
- [0183] Pechan, P.M., Bartels, D., Brown, D.C.W. and Schell, J. (1991). Messenger-RNA and protein changes associated with induction of *Brassica microspore* embryogenesis. *Planta* 184, 161-165.
- [0184] Roberts, M.R., Foster, G.D., Blundell, R.P., Robinson, S.W., Kumar, A., Draper, J. and Scott, R. (1993). Gametophytic and sporophytic expression of an anther-specific *Arabidopsis thaliana* gene. *Plant J.* 3, 111-120.
- [0185] Rounsley, S.D., Ditta, G.S. and Yanofsky, M.F. (1995). Diverse roles for MADS box genes in *Arabidopsis* development. *Plant Cell* 7, 1259-1269.
- [0186] Salter, M.G., Paine, J.A., Riddell, K.V., Jepson, I., Greenland, A.J., Caddick, M.X., Tomsett, A.B. (1998). Characterisation of the ethanol-inducible alc gene expression system for transgenic plants. *Plant Journal* 16, 127-132.
- [0187] Sambrook, J., Fritsch, E.F. and Maniatis, T. (1989). *Molecular Cloning: A Laboratory Manual*, second edition. (Cold Spring Harbor: Cold Spring Harbor Laboratory Press).
- [0188] Schmidt, E.D.L., Guzzo, F., Toonen, M.A.J. and de Vries, S.C. (1997). A leucine-rich repeat containing receptor-like kinase marks somatic plant cells competent to form embryos. *Development*. 124, 2049-2062.
- [0189] Stockinger, E.J., Gilmour, S.J. and Thomashow, M.F. (1997). *Arabidopsis thaliana* CBF1 encodes an AP2 domain-containing transcriptional activator that binds to the C-repeat/DRE, a cis-acting DNA regulatory element that stimulates transcription in response to low temperature and water deficit. *Proc. Natl. Acad. Sci. USA* 94, 1035-1040.
- [0190] Taylor, R.L. (1967). The foliar embryos of *Malaxis paludosa*. *Can. J. Bot.* 45, 1553-1556.
- [0191] Treacy, B.K., Hattori, J., Prud'homme, I., Barbour, E., Boutilier, K., Baszczyński, C.L., Huang, B., Johnson, D.A. and Miki, B.L. (1997). Bnml, a *Brassica* pollen-specific gene. *Plant Mol. Biol.* 34, 603-611.
- [0192] Twell, D., Wing, R., Yamaguchi, J. and McCormick, S. (1989). Isolation and expression of an anther-specific gene from tomato. *Mol. Gen. Genet.* 217 240-245.
- [0193] Ulmasov, T., Murfett, J., Hagen, G. and Guilfoyle, T. (1997). Aux/IAA proteins repress expression of reporter genes containing natural and highly active synthetic auxin response elements. *Plant Cell* 9, 1963-1971.
- [0194] van Engelen, F.A., Molthoff, J.W., Conner, A.J., Nap, J.P., Pereira, A. and Stiekema, W.J. (1995). pBINPLUS: an improved plant transformation vector based on pBIN19. *Transgenic Res.* 4, 288-290.
- [0195] Vergunst, A.C., de Waal, E.C. and Hooykaas, P.J.J. (1998). Root transformation by *Agrobacterium tumefaciens*. In *Arabidopsis Protocols*, J.M. Martinez-Zapater and J. Salinas, eds (Totowa, USA: Humana Press).
- [0196] Xu, D., McElroy, D., Thomburg, R.W. and Wu, R.C.S. (1993). Systemic induction of a potato pin⁺ promoter by wounding, methyl jasmonate, and abscisic acid in transgenic rice plants. *Plant Mol. Biol.* 22, 573-588.
- [0197] Xu, H., Knox, R.B., Taylor, P.E. and Singh, M.B. (1995). Bcpl, a gene required for male fertility in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* 92, 2106-2110.
- [0198] Yarbrough, J.A. (1932). Anatomical and developmental studies of the foliar embryos of *Bryophyllum calycinum*. *Amer. J. Bot.* 19, 443-453.
- [0199] Yeung, E.C. (1995). Structural and developmental patterns in somatic embryogenesis. In *In Vitro Embryogenesis in Plants*. T.A. Thorpe, ed (Dordrecht: Kluwer Academic Publishers), pp. 205-247.
- [0200] Zarsky, V., Garrido, D., Eller, N., Tupy, J., Vicente, O., Schaffl, F. and Heberle-Bors, E. (1995). The expression of a small heat shock gene is activated during induction of tobacco pollen embryogenesis by starvation. *Plant Cell Environ.* 18, 139-147.
- [0201] Zhou, J., Tang, X. and Martin, G.B. (1997). The Pto kinase conferring resistance to tomato bacterial speck disease interacts with proteins that bind a cis-element of pathogenesis-related genes. *EMBO J.* 16, 3207-3218.

SEQUENCE LISTING

5 <110> CPRO-DLO
Agriculture and Agri-food Canada

<120> Use of the BNM3 Transcriptional Activator to Control
10 Plant Embryogenesis and Regeneration Processes

<130> E158138

<140>
15 <141>

<160> 12

<170> PatentIn Ver. 2.1

<210> 1
<211> 2014
<212> DNA
25 <213> Brassica napus

<400> 1
gttcacatctct cttctttaag accaaaacct tttctctctc ctcttcacgc atgaacccta 60
30 actaagttctt tctcttttta ctttttacca agaactcggt agatcactct ctgaactcaa 120
tgaataataaa ctgggttaggc tttctctctc ctctctatga acaaaatcac catcgtaagg 180
acgtctactc ttcaccacc acaaccgtcg tagatgtcgc cggagagtac tgttacgac 240
cgacccgtgc ctccgatgag tcttcagcca tccaaacatc gttctctctc cctcttggtg 300
tcgtcgtcga tgccttcacc agagacaaca atagtcactc ccgagattgg gacatcaatg 360
35 gttgtgcgat caataacatc cacaacgatg agcaagatgg accaaagctt gagaatttcc 420
ttggccgcac caccacgatt tacaacacca acgaaaaact tggagatgga agtggaaagt 480
gctgttatgg aggaggagac ggtggtggtg gctcactagg actttcgatg ataaagacat 540
ggctgagaaa tcaacccgtg gataatggtg ataatacaaga aaatggcaat gctgcataag 600
gctctgctct ctcaatgaac tctctactt cttgtgataa caacaacgac agcaataaca 660
40 acgtgtgttc ccaagggaag actattgatg atagcgttga agctacaccg aagaaaaacta 720
ttgagagatt tggacagagg acgtctatat acccggtgtg tacaaggcat cgggtggacg 780
gaagatatga ggcacattta tgggataata gttgtaaaag agaagggcaa acgcgcacaa 840
45 aaagacaagt ttatttggga ggttatgaca aagaagaaaa agcagctagg gctttatgatt 900
tagccgcact caagtatgg ggaaccacca ctactactaa cttccccatg agcgaatatg 960
aaaaagaggt agaagagatg aaacacatga caaggcaaga gtaagtgtcc tcaatcgcca 1020
ggaaaaagtag tggttctctc cgtggtgcat cgatttatcg tggagtaaca agacatcacc 1080
aacatggaaag atggcaagct aggataggaa gactcgccgg taacaaagac ctctacttgg 1140
50 gaacttttgg caccacaagaa gaagctgcag aggcatacga catttgcggcc atcaaaattca 1200
gaggattaac ccgactgact aacttcgaca tgaacagata caacgttaaa gcaatctctg 1260
aaagccctag tcttctctatt ggtagcgccg caaaaagctc caaggaggtc aacgttccg 1320
ttccaagtaa gatgatgac agtaataacg ttccagagag tgagaaatagt gctagcgggt 1380
ggcnaaacgc tgcggttcag catcatcagg gactagattt gactttattg caccacatc 1440
55 aaagagagta caatcggtat tattacaatg gaggaaactt gtcttcgag agtgcaggag 1500

cttgtttcaa acaagaggat gatcaacacc atttcttgag caacacgcag agcctcatga 1560
 ctaatatcga tcatcaaaagt tctgtttcgg atgattcggg tactgtttgt ggaaatgttg 1620
 5 ttgggttatgg tgggttatcaa ggatttgcag ccccggttaa ctgcgaatgcc tacgtctgta 1680
 gtgagtttga ttataacgca agaaaccatt attactttgc tcagcagcag cagacccagc 1740
 agtcgccagg tggagatttt cccgcggcaa tgacgaataa tgttggctct aatatgtatt 1800
 accatgggga aggtgggtga gaagttgctc caacatttac agtttggaaac gacaaataga 1860
 10 aaaaatagtt aaagatcttt agttatatgc gttgttgtgt gctgggtaac agtgtgatac 1920
 tttgattatg tttttttctt tctcttttct tttttcttgg ttaatttctt aagaacttatt 1980
 tttagtttcc attagttaga taaattttca gact 2014

15 <210> 2
 <211> 579
 <212> PRT
 <213> Brassica napus

20 <400> 2
 Met Asn Asn Asn Trp Leu Gly Phe Ser Leu Ser Pro Tyr Glu Gln Asn
 1 5 10 15

25 His His Arg Lys Asp Val Tyr Ser Ser Thr Thr Thr Thr Val Val Asp
 20 25 30

Val Ala Gly Glu Tyr Cys Tyr Asp Pro Thr Ala Ala Ser Asp Glu Ser
 30 35 40 45

Ser Ala Ile Gln Thr Ser Phe Pro Ser Pro Phe Gly Val Val Val Asp
 50 55 60

35 Ala Phe Thr Arg Asp Asn Asn Ser His Ser Arg Asp Trp Asp Ile Asn
 65 70 75 80

Gly Cys Ala Cys Asn Asn Ile His Asn Asp Glu Gln Asp Gly Pro Lys
 40 85 90 95

Leu Glu Asn Phe Leu Gly Arg Thr Thr Thr Ile Tyr Asn Thr Asn Glu
 100 105 110

45 Asn Val Gly Asp Gly Ser Gly Ser Gly Cys Tyr Gly Gly Gly Asp Gly
 115 120 125

Gly Gly Gly Ser Leu Gly Leu Ser Met Ile Lys Thr Trp Leu Arg Asn
 50 130 135 140

Gln Pro Val Asp Asn Val Asp Asn Gln Glu Asn Gly Asn Ala Ala Lys
 145 150 155 160

55 Gly Leu Ser Leu Ser Met Asn Ser Ser Thr Ser Cys Asp Asn Asn Asn

	163	170	175
5	Asp Ser Asn Asn Asn Val Val Ala Gln Gly Lys Thr Ile Asp Asp Ser 180	185	190
10	Val Glu Ala Thr Pro Lys Lys Thr Ile Glu Ser Phe Gly Gln Arg Thr 195	200	205
	Ser Ile Tyr Arg Gly Val Thr Arg His Arg Trp Thr Gly Arg Tyr Glu 210	215	220
15	Ala His Leu Trp Asp Asn Ser Cys Lys Arg Glu Gly Gln Thr Arg Lys 225	230	235 240
20	Gly Arg Gln Val Tyr Leu Gly Gly Tyr Asp Lys Glu Glu Lys Ala Ala 245	250	255
	Arg Ala Tyr Asp Leu Ala Ala Leu Lys Tyr Trp Gly Thr Thr Thr Thr 260	265	270
25	Thr Asn Phe Pro Met Ser Glu Tyr Glu Lys Glu Val Glu Glu Met Lys 275	280	285
30	His Met Thr Arg Gln Glu Tyr Val Ala Ser Leu Arg Arg Lys Ser Ser 290	295	300
	Gly Phe Ser Arg Gly Ala Ser Ile Tyr Arg Gly Val Thr Arg His His 305	310	315 320
35	Gln His Gly Arg Trp Gln Ala Arg Ile Gly Arg Val Ala Gly Asn Lys 325	330	335
40	Asp Leu Tyr Leu Gly Thr Phe Gly Thr Gln Glu Glu Ala Ala Glu Ala 340	345	350
	Tyr Asp Ile Ala Ala Ile Lys Phe Arg Gly Leu Thr Ala Val Thr Asn 355	360	365
45	Phe Asp Met Asn Arg Tyr Asn Val Lys Ala Ile Leu Glu Ser Pro Ser 370	375	380
50	Leu Pro Ile Gly Ser Ala Ala Lys Arg Leu Lys Glu Ala Asn Arg Pro 385	390	395 400
	Val Pro Ser Met Met Met Ile Ser Asn Asn Val Ser Glu Ser Glu Asn 405	410	415
55	Ser Ala Ser Gly Trp Gln Asn Ala Ala Val Gln His His Gln Gly Val		

	420	425	430
5	Asp Leu Ser Leu Leu His Gln His Gln Glu Arg Tyr Asn Gly Tyr Tyr 435 440 445		
10	Tyr Asn Gly Gly Asn Leu Ser Ser Glu Ser Ala Arg Ala Cys Phe Lys 450 455 460		
	Gln Glu Asp Asp Gln His His Phe Leu Ser Asn Thr Gln Ser Leu Met 465 470 475 480		
15	Thr Asn Ile Asp His Gln Ser Ser Val Ser Asp Asp Ser Val Thr Val 485 490 495		
20	Cys Gly Asn Val Val Gly Tyr Gly Gly Tyr Gln Gly Phe Ala Ala Pro 500 505 510		
	Val Asn Cys Asp Ala Tyr Ala Ala Ser Glu Phe Asp Tyr Asn Ala Arg 515 520 525		
25	Asn His Tyr Tyr Phe Ala Gln Gln Gln Gln Thr Gln Gln Ser Pro Gly 530 535 540		
30	Gly Asp Phe Pro Ala Ala Met Thr Asn Asn Val Gly Ser Asn Met Tyr 545 550 555 560		
	Tyr His Gly Glu Gly Gly Gly Glu Val Ala Pro Thr Phe Thr Val Trp 565 570 575		
35	Asn Asp Asn		
40	<210> 3 <211> 2011 <212> DNA <213> Brassica napus		
45	<400> 3 ttcttctttt accctttacc aagaactcgt tagatcattt tctgaactcg atgaataata 60 actgggttagg cttttctctc tctccttatg aacaaaatca ccactgtaag gacgtctgct 120 cttccaccac cacaaccgcc gtagatgctg ccggagagta ctgttaacgat ccgaccgctg 180 cctccgatga gtcttcagcc atccaaacat cgttctcttc tccctttggt gtcgtctctg 240 atgctttcac cagagacaac aatagtcact ccggagattg ggacatcaat ggtagtccat 300 gtaataacat ccacaatgat gagcaagatg gacaaaaact tgagaatttc ctggcgccga 360 ccaccacgat ttacaacacc aacgaaaacg ttggagatac cgatggaagt ggggtgttatg 420 gaggaggaga cgggtgggtggg ggctcactag gaatttcgat gataaagana tggctgagaa 480		

atcaaccggt ggataatgtt gataatcaag aaaatggcaa tgggtgcaaaa ggctgtgtcc 540
 tctcaatgaa ctcatctact tcttgtgata acaacaacta cagcagtaac aaacctgttg 630
 5 cccaagggaa gactattgat gatagcggtt aagctacacc gaagaaaaat attgagagtt 660
 ttggcagag gagctctata taccgcggtg ttacaaggca tcggtggaca ggaagatatg 720
 aggcacattt atgggataat agttgtaaac gagaaggcca aacgcgcgaaa ggaagacaag 780
 ttattttggg aggttatgac aaagaagaaa aagcagctag ggcttatgat ttagccgcac 840
 10 tcaagtattg gggaaccacc actactacta acttcccat gagcgaatat gagaagaga 900
 tagaaagagat gaagcacatg acaaggcaag agtatgttc ctcacttcgc aggaaaagta 960
 gtggtttctc tcgtggtgca tcgattatc gtggagtaac aagacatcac caacatggaa 1020
 gatggcaagc taggatagga agagtgcgcg gtaacaaaaga cctcacttg ggaacttttg 1080
 gcacacaaga agaagctgca gaggcatacg acattgcggc catcaaatc aggaggattaa 1140
 15 ccgcagtgac taacttcgac atgaacagat acaacgttaa agcaatctc gaaagcccta 1200
 gtcttcctat tggtagcgcc gcaaaacgct tcaacggagg taaccgtccg gttccaagta 1260
 tgatgatgat cagtaataac gtttcagaga gtgaacaataa tgctagcggt tggcaaaacg 1320
 ctgcggttca gcatcatcag ggagttagatt tgagcttatt gcagcaaat caagagaggt 1380
 20 acaatcggtta ttattacaat ggagaaaact tgtcttcgga gagtgttagt gcttgtttca 1440
 aacaagagga tgatcaacac catttcttga gcaacacgca gagctctatg actaatatcg 1500
 atcatcaaag tctgttttca gatgattcgg ttaactgttg tggaaatgtt gttggttatg 1560
 gtggtttatca aggattttgca gccccggtta actgcgatgc ctacgctgt agtgagtttg 1620
 25 actataacgc aagaaacat tattactttg ctacgacgca gcagaccag cattcgccag 1680
 gaggagattt tcccgcggca atgacgaata atgttggtc taatatgtat taacatgggg 1740
 aagtggttg agaatgtgct ccaacattta cagtttgga cgacaattg aaataatagt 1800
 taaagatctt tagttatatg cgttgttgtg ttgtgttgaa cagtttgata ctttgattat 1860
 gttttttttt ctctttttca ttttgttgtg tagtttctta agacttattt ttgttttcca 1920
 30 ttagtggat aaattttcgg acttaagggt cacttctgtt ctgactcttg tctaatacag 1980
 aaaagttttc ataaaaaaaa aaaaaaaaaa a 2011

<210> 4
 35 <211> 579
 <212> PRT
 <213> Brassica napus

 40 <400> 4
 Met Asn Asn Asn Trp Leu Gly Phe Ser Leu Ser Pro Tyr Glu Gln Asn
 1 5 10 15

 45 His His Arg Lys Asp Val Tyr Ser Ser Thr Thr Thr Val Val Asp
 20 25 30

 Val Ala Gly Glu Tyr Cys Tyr Asp Pro Thr Ala Ala Ser Asp Glu Ser
 35 40 45

 50 Ser Ala Ile Gln Thr Ser Phe Pro Ser Pro Phe Gly Val Val Val Asp
 50 55 60

 55 Ala Phe Thr Arg Asp Asn Asn Ser His Ser Arg Asp Trp Asp Ile Asn
 65 70 75 80

29

Asp Leu Tyr Leu Gly Thr Phe Gly Thr Gln Glu Glu Ala Ala Glu Ala
 5 340 345 350
 Tyr Asp Ile Ala Ala Ile Lys Phe Arg Gly Leu Thr Ala Val Thr Asn
 355 360 365
 10 Phe Asp Met Asn Arg Tyr Asn Val Lys Ala Ile Leu Glu Ser Pro Ser
 370 375 380
 Leu Pro Ile Gly Ser Ala Ala Lys Arg Leu Lys Glu Ala Asn Arg Pro
 15 385 390 395 400
 Val Pro Ser Met Met Met Ile Ser Asn Asn Val Ser Glu Ser Glu Asn
 405 410 415
 20 Ser Ala Ser Gly Trp Gln Asn Ala Ala Val Gln His His Gln Gly Val
 420 425 430
 Asp Leu Ser Leu Leu His Gln His Gln Glu Arg Tyr Asn Gly Tyr Tyr
 25 435 440 445
 Tyr Asn Gly Gly Asn Leu Ser Ser Glu Ser Ala Arg Ala Cys Phe Lys
 450 455 460
 30 Gln Glu Asp Asp Gln His His Phe Leu Ser Asn Thr Gln Ser Leu Met
 465 470 475 480
 Thr Asn Ile Asp His Gln Ser Ser Val Ser Asp Asp Ser Val Thr Val
 35 485 490 495
 Cys Gly Asn Val Val Gly Tyr Gly Gly Tyr Gln Gly Phe Ala Ala Pro
 500 505 510
 40 Val Asn Cys Asp Ala Tyr Ala Ala Ser Glu Phe Asp Tyr Asn Ala Arg
 515 520 525
 Asn His Tyr Tyr Phe Ala Gln Gln Gln Gln Thr Gln Gln Ser Pro Gly
 45 530 535 540
 Gly Asp Phe Pro Ala Ala Met Thr Asn Asn Val Gly Ser Asn Met Tyr
 545 550 555 560
 50 Tyr His Gly Glu Gly Gly Gly Glu Val Ala Pro Thr Phe Thr Val Trp
 565 570 575
 55 Asn Asp Asn

```

5
    <210> 5
    <211> 4873
    <212> DNA
    <213> Brassica napus
10
    <220>
    <221> intron
    <222> [1846]..(2298)
15
    <220>
    <221> intron
    <222> [2720]..(2952)
20
    <220>
    <221> intron
    <222> [3036]..(3160)
25
    <220>
    <221> intron
    <222> [3170]..(3314)
30
    <220>
    <221> intron
    <222> [3404]..(3553)
35
    <220>
    <221> intron
    <222> [3623]..(3797)
40
    <220>
    <221> intron
    <222> [3849]..(3961)
45
    <220>
    <221> intron
    <222> [4039]..(4148)
50
    <220>
    <221> misc_feature
    <222> [1620]..(1622)
    <223> start codon
55
    <220>
    <221> misc_feature
    <222> [4856]..(4858)

```

<223> stop codon

5 <400> 5
 atctctccac cgattcgtta cccagtgctt gaaaatatga tgcctacgaa tcaattaaat 60
 ggagaagctc cactgcttgt gtaggtagaa gctcaagcaa caaccgaaa cctcggcgtt 120
 atcgggaggt agcatcgtta ttgcacaaaa ttccgcgcgc agagatgaaa cgattcaaga 180
 10 gaaaccccca aataggttag ccataaaaca gtgaattagt atgatttaag agataaagaag 240
 agaagatgag ttcaagaaaa gaaataccca catctattta tactgtttac acaccgcctt 300
 tcaagtctaa gcaaaagcatt gaagatgaat cgtggaggag agttaaatag atttaacaca 360
 aagccatttaa ccaaacccgtt gcaggtcggg agacgaaccc caaaagtcac gcttagccgt 420
 cgcacgaaga ggagcgatga atttcgtttt ctcgctgcag tcgtattagg gatagacgga 480
 15 gctcattatc gttgggcggg aaacacttct aatctcaacg cccatgaaca cactaaagaa 540
 cgaaaccgaa aatgtttgaa gtttaatgaa acgtgcgggtt tgccttatgg acacatgtca 600
 ttacgatata aatgatatta tctacgtgga tcataggtgt ctctctaagg agagagcaaa 660
 cctatacttt atataaatag atttgatca ttctaaggag tgtttaagat ttctgataa 720
 20 atattaaaaa aaattacaaa tttttatgta attagttttg ttacataaaa ataactaaa 780
 ataaaaataa tcaaccaaat aaaaaaacac ggtattttat aattggtcaa aaaaaaaaat 840
 aaaaacttaa atttcacctt gaattacgag aatgtcaact attttgaaac aaactcaaaa 900
 tctttaaaca tcaattaaac tgatacggat ggagtatata tctttacaga gaacatatat 960
 atatgttttt ctgttaagcg tccatctctt cttagtcatg tagttcaaat accagctgca 1020
 25 gtaaaaccat gaattattga atttgtgtga aaatatccga agcgactact gcaagtttgg 1080
 aagcaaaacg ccaaacgcaa tcgctcgcctt ggctataggg tcaacacata acatgtgact 1140
 agcatattgg gtcttaattc aacagcgagt gattttggga ttactatta gttcctgtgt 1200
 tactctactt ttaacacaaa gtcaactaac ttatttacac atgaagagag gtttgaagag 1260
 30 gcttttgact gattaattat aatgtattaa accaaactag aattaaagaa ttaggctatt 1320
 aattacatta ccaccaccac ccaccattca acccgaccaa tcaactcca caggtttcaa 1380
 gtaaaacaa ttttttttgt tgttctctcg gaattttaat aaatatctgt ttatataaat 1440
 gcgcagtata tgagcgcctc gaagaaatga aacattatat ctttgacttt tctctcctta 1500
 gttcctctct cttctttaag accaaaacct tttctctctc ctctctcatg atgaacctta 1560
 35 actaagttct tcttctttta ccttttacc aagaactcgt agatcactct ctgaactcaa 1620
 tgaataataa ctggttaggc ttttctctct ctctctatga acaaaatcac catcgtaagg 1680
 acgtctactc ttccaccacc acaacgcgtc tagatgtcgc cggagagtag tgttaccgat 1740
 cgaccgcgtc ctccgatgag tcttcagcca tccaaacatc gtttctctct ccttttgggt 1800
 40 tcgtcgtcga tgccttccac agagacacaa atagctactc ccgaggttat tgttttagaa 1860
 ctacttgctt ttttttgatt tgtttatttg tttagtttcc tttcttcca atcgtagaaa 1920
 caaagaccaa taccacgcga cgcatactag ccttattttt tctttgggct tatttatcga 1980
 tttcatttat ttgagaata tcaatgtgtg ggttttgatg ttgtttgca tatagtaata 2040
 45 ctaaaacata tgcaggttat acatagattt ttttttaaga tatacatgga tatgaaatga 2100
 aatttgacat tctctctctt attcaatcac ataattgat cacatacatg tgcacctttt 2160
 gatttgtata ttgttttctt acagttgaag gagagaataa ccaaaatccc atttgatat 2220
 tatagatcgg tgatgaaaag taaatttaac aaattatgat aatatagccc attaatcttt 2280
 gatctttttt ctttatagat tgggacatca alggtttgcg atgcaataac atcaacaacg 2340
 50 atgagcaaga tggaccaaa gttgagaatt tctttggcgc caccaccacg atttacaaca 2400
 ccaacgaaaa cgtttggagat ggaagtggaa gtggctgtaa tggaggagga gacgttgggt 2460
 gtggctcact aggacttttc atgataaaga catggctgag aaatacaacc gtggaataatg 2520
 ttgataatca agaaaaatggc aatgctgcaa aaggcctgtc cctctcaaat aactcaatca 2580
 55 cttcttgcga taacaacaa gacagcaata caaacgtgt tgccttaggg aagactatcg 2640
 atgatacgtc tgaagctaca ccgaagaaaa ctattgagag ttttgacag aggcagctca 2700

tataccgogg tgttacaagg tgccttcat ttattttaat aaaaagtga aaatgtcgct 2760
 tgaattgtta tcttcttggg aaagctggg acattgatct aatggctctg ttgcgagagt 2820
 5 gctaccgaat ggtccttgat atatatgato aaagagagat attgttatta tgggcttata 2880
 tagaataata catatatata tatatatca tggtagctgt tgatgacatg tatgttccta 2940
 ttaaatgata aggcattcgg ggacagggaag atatgaggca catttacggg ataatagttg 3000
 taaaagagaa ggccaaacgc gcaaagggaag acaaggatata tatataacca ttgataaatt 3060
 10 gatcataatt tcatacacga ttactttca aactaaataa ggtttttcga tcaattgttca 3120
 tgtttttatc aaaaatttga cctgttggtt gtcttctcag ttatttggg taagttaatt 3180
 attataaatt ggacgaagct gtgatgggta aatctaaatt atataatcaa atttgtttat 3240
 tttttggta taccattcat atataatcaa aatagcgata cgaattctac tcaattgttg 3300
 tctatatcat gcaggagggt atgacaaaga agaaaaagca gctagggcct atgatttagc 3360
 15 cgactcaag tatitgggga ccaccatcac tactaacctc ccgtaagtc aatcaatgtt 3420
 gtacaagatt tcataactta gaaccaattt tattcttttt ttataagatg ctattatctt 3480
 attttaaatt gccatgttta tatcgttaca ttattaccaa taaaagtac ttitgggttg 3540
 atataatagt tagatgagcg aatatgaana agaggtagaa gagatgaagc acatgacnag 3600
 20 gcaagagtat gttgcctcac tgcgcaggta tataatggaa ctctcgatat tattgcatat 3660
 ggcatttatt attatcatg tatattagta ttttatata agaaccatc acgctcacgt 3720
 ttntatttaa naatatgtcc gtattcacgt cagattatca gcataccct atatatata 3780
 gacattaaaa tatgcaggaa agtagtggt ttctctgtg gtgcacgat ttatcgttga 3840
 gtaaccaagg attcatcacg agagaacgaa tctattttg ttacgtacat atatatataa 3900
 25 aaataaatt ataagatatc acattttata ttatgaataa ttcttctaatt gggtccaaaa 3960
 gacatcacca acatggaaga tggcaagcta ggatagggaag agtcgcgggt acaaaagacc 4020
 tctacttggg aacttttggg acgttttact ttcttctact aaacttcaca atcaaatcta 4080
 taacaaaaga tatcaactaa aaactacaac atatatctaa gtaagctgta catatattat 4140
 30 atatgaagg acacaagaag aagctgcaga ggcatcacg attgcggcca tcaaatcagc 4200
 aggatcaacc gcagtgcata acttcgacat gaacagatc aacgttaag caatcctcga 4260
 aagccctagt ctctctattg gtacgcgcgc aaaaagctctc aaggaggcta accgtccggg 4320
 tccaagtatg atgatgatca gtaataacgt ttcatgagat gagaatagtg ctacgggttg 4380
 35 gcaaaacgct gcggttcagc atcatcaggg agtagatttg agcttattgc accaactatca 4440
 agagaggtaac aatgggttatt attacaatgg aggaacattg tcttcggaga gtgtagggc 4500
 ttgtttcaaa caagaggatg atcaaacacca ttctctgagc aacacgcaga gctcatgac 4560
 taatatcgat catcaaatgt ctgtttcggg tgatcgggtt actgtttgtg gaaatgttg 4620
 40 tggttatggt ggttatcaag gatttgcagc ccggttaac tgcgatgct acgctgctag 4680
 tgagtttgat tataacgcaa gaacacatta ttactttgtc cagcagcagc agaccacga 4740
 gtgcacaggt ggagattttc ccgcggaat gacgaataat gttggtctta atatgtatta 4800
 ccattggggaa ggtgttggag aagttgtctc aacatttaca gtttggaaag acaattagaa 4860
 aaaaatagtta aag 4873

<210> 6

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 6

gaggcagcgg tcggatcgta acagtactct

30

5

<210> 7

<211> 30

<212> DNA

<213> Artificial Sequence

10

<220>

<223> Description of Artificial Sequence:primer

15

<400> 7

cataaaggaga gagagaaaag cctaaccagt

30

20

<210> 8

<211> 19

<212> DNA

<213> Artificial Sequence

25

<220>

<223> Description of Artificial Sequence:primer

<400> 8

accaagaact cgtagatc

30

19

<210> 9

<211> 20

<212> DNA

<213> Artificial Sequence

35

<220>

<223> Description of Artificial Sequence:primer

40

<400> 9

aacgcacata actaaagatc

20

45

<210> 10

<211> 26

<212> DNA

<213> Artificial Sequence

50

<220>

<223> Description of Artificial Sequence:primer

55

<400> 10

ccatggatcc agagacgaag cgaaac

26

5

<210> 11

<211> 26

<212> DNA

10

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

15

<400> 11

actccatgga taataactgg ttaggc

26

20

<210> 12

<211> 26

<212> DNA

<213> Artificial Sequence

25

<220>

<223> Description of Artificial Sequence:primer

30

<400> 12

aaattctcaa gctttggtcc atcttg

26

35

40

45

50

55

Claims

1. An isolated DNA molecule comprising a nucleotide sequence that hybridizes to SEQ ID NO:5 under stringent conditions
2. The isolated DNA molecule of claim 1 wherein said isolated DNA molecule comprises at least 23 contiguous nucleotides of SEQ ID NO:5.
3. The isolated DNA molecule of claim 1 wherein said isolated DNA molecule comprises a nucleotide sequence that is at least 70% homologous with the nucleotide sequence defined by SEQ ID NO:5.
4. An isolated DNA molecule comprising a nucleic acid sequence encoding a protein, wherein said protein when present at a sufficient level within a plant cell renders said cell embryogenic, increases the regenerative capacity of said plant cell, or both renders said plant cell embryogenic and increases the regenerative capacity of said plant cell, said isolated DNA molecule having at least 70% homology within a nucleotide sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3 and SEQ ID NO:5.
5. The isolated DNA molecule of claim 4 comprising a nucleotide sequence that hybridizes to nucleotides 1620-4373 of SEQ ID NO:5 under stringent conditions.
6. The isolated DNA molecule of claim 4 comprising a nucleotide sequence that hybridizes to the nucleotide sequence of SEQ ID NO:1 under stringent conditions.
7. The isolated DNA molecule of claim 4 comprising a nucleotide sequence that hybridizes to the nucleotide sequence of SEQ ID NO:3 under stringent conditions.
8. The isolated DNA molecule of claim 6, wherein said DNA encodes a protein as defined by SEQ ID NO:2.
9. The isolated DNA molecule of claim 7, wherein said DNA encodes a protein as defined by SEQ ID NO:4.
10. A vector comprising the isolated DNA molecule as claimed in any one of claims 1 to 9, wherein said isolated DNA molecule is under control of a regulatory element that directs expression of said DNA in a plant cell.
11. The vector of claim 10, wherein said regulatory element is a constitutive regulatory element
12. The vector of claim 10, wherein said regulatory element is an inducible regulatory element.
13. The vector of claim 10, wherein said regulatory element is a tissue specific regulatory element
14. The vector of claim 10, wherein said regulatory element is an developmentally active regulatory element.
15. A transformed plant cell comprising the vector of any one of claims 10 to 14.
16. A transformed plant comprising the vector of any one of claims 10 to 14.
17. A seed obtained from the transformed plant of claim 16.
18. An isolated protein encoded by the isolated DNA molecule as claimed in any one of claims 4 to 9.
19. A method of producing asexually derived embryos comprising:
 - i) transforming a plant cell with the vector of any one of claims 10 to 14;
 - ii) growing said plant cell to produce transformed tissue;
 - iii) selecting said transformed tissue for occurrence of said isolated DNA molecule; and
 - iv) assaying said transformed plant for asexual embryo production.
20. The method of claim 19 wherein the step of assaying involves assaying for adventitious embryony.

21. The method of claim 19, wherein the step of assaying involves assaying for somatic embryos
22. The method of claim 19, wherein the step of assaying involves assaying for gametophytic embryos.
- 5 23. The method of claim 19, wherein the step of assaying involves assaying for haploid parthenogenesis of the embryo sac.
24. The method of claim 19, wherein the step of assaying involves assaying for diplospory.
- 10 25. A method of modifying the regenerative capacity of a plant comprising
 - i) transforming a plant cell with the vector of any one of claims 10 to 14;
 - ii) growing said transformed plant cell to produce transformed tissue; and
 - 15 iii) assaying said transformed plant tissue for enhanced regeneration as compared to wild-type tissue.
26. The method of claim 25, wherein the step of growing said transformed plant cell, the step of assaying said transformed plant tissue, or both the step of growing said transformed plant cell and the step of assaying said transformed plant tissue are carried out in the absence of a growth regulator.
- 20 27. A method of selecting a transformed plant comprising:
 - i) transforming a normally non-regenerative plant with a vector of any one of claims 10 to 14; and
 - 25 ii) determining whether said transformed plant is able to regenerate under conditions in which said normally non-regenerative plant does not regenerate.
28. The isolated DNA molecule of claim 1 comprising a DNA sequence that hybridizes to nucleotides 1-1619 of SEQ ID NO:5 under stringent conditions.
29. The isolated DNA molecule of claim 1 wherein said isolated molecule comprises at least 22 contiguous nucleotides within nucleotides 1-1619 of SEQ ID NO:5.
- 30 30. A vector comprising the isolated DNA molecule of either claim 28 or 29 operably associated with a gene of interest, wherein said isolated DNA molecule directs the expression of said gene of interest within a plant cell.
- 35 31. The vector as defined by claim 30, wherein said gene of interest is heterologous with respect to the isolated DNA molecule.
32. The vector as defined by claim 31, wherein said gene of interest is selected from the group consisting of a pharmaceutically active protein, antibody, industrial enzyme, protein supplement, nutraceutical, storage protein, animal feed and animal feed supplement.
- 40 33. A transformed plant cell comprising the vector of either claim 30, 31 or 32.
34. A transformed plant comprising the vector of either claim 30, 31 or 32.
- 45 35. A seed obtained from the transformed plant of claim 34.
36. A method for directing the expression of a gene of interest within a developing embryo of a plant comprising transforming said plant with the vector as defined by either claim 30, 31 or 32.
- 50 37. A use of a nucleotide sequence as defined in any one of claims 4, 5, 6 or 7 as a selectable marker.
38. A method of producing asexually derived embryos comprising:
 - 55 i) transiently transforming a plant cell with the vector of any one of claims 10 to 14, or introducing into said plant cell the protein of claim 18, to produce a modified plant cell;
 - ii) growing said modified plant cell to produce tissue; and
 - iii) assaying said tissue for asexual embryo formation.

39. The method of claim 38 wherein the step of assaying involves assaying for adventitious embryony.

40. The method of claim 38, wherein the step of assaying involves assaying for somatic embryos.

41. The method of claim 38, wherein the step of assaying involves assaying for gametophytic embryos

42. The method of claim 38, wherein the step of assaying involves assaying for haploid parthenogenesis of the embryo sac.

43. The method of claim 38, wherein the step of assaying involves assaying for diplospory.

44. A method of modifying the regenerative capacity of a plant comprising

- i) transiently transforming a plant cell with the vector of any one of claims 10 to 14 or introducing into said plant cell the protein of claim 18, to produce a modified plant cell;
- ii) growing said modified plant cell to produce tissue; and
- iii) assaying said tissue for enhanced regeneration as compared to wild-type tissue.

45. The method of claim 44, wherein the step of growing said modified plant cell, the step of assaying said tissue, or both the step of growing said modified plant cell and the step of assaying said tissue are carried out in the absence of a growth regulator.

46. A method of producing an apomictic plant comprising:

- i) transforming a plant with the vector of any one of claims 10 to 14, to produce a transformed plant;
- ii) selecting said transformed plant for occurrence of said isolated DNA molecule; and
- iii) assaying said transformed plant for asexual embryo production.

47. The method of claim 46 wherein the step of assaying involves assaying for adventitious embryony.

48. The method of claim 46, wherein the step of assaying involves assaying for somatic embryos.

49. The method of claim 46, wherein the step of assaying involves assaying for gametophytic embryos.

50. The method of claim 46, wherein the step of assaying involves assaying for parthenogenesis of the embryo sac.

51. A method of modifying the regenerative capacity of a plant comprising

- i) transiently transforming a plant cell with the vector of any one of claims 10 to 14, or introducing into said plant cell the protein of claim 18;
- ii) growing said plant cell to form tissue; and
- iii) assaying said tissue for enhanced regeneration as compared to wild-type tissue.

52. The method of claim 51, wherein the step of growing said plant cell, the step of assaying said tissue, or both the step of growing said plant cell and the step of assaying said tissue are carried out in the absence of a growth regulator.

53. A method of selecting a modified plant comprising:

- i) transiently transforming a normally non-regenerative plant with a vector of any one of claims 10 to 14, or introducing into said normally non-regenerative plant the protein of claim 18, to produce said modified plant; and
- ii) determining whether said modified plant is able to regenerate under conditions in which said normally non-regenerative plant does not germinate.

54. An isolated DNA molecule comprising a sequence encoding a protein consisting of two AP2 DNA binding domains, which when said protein is expressed at a sufficient level in a plant cell, renders said cell embryogenic, or increase the regenerative capacity of said plant cell, or both renders said cell embryogenic and increase the regenerative

capacity of said plant cell.

55. A method of producing a protein of interest comprising

- i) transforming a plant with at least one vector, said at least one vector selected from any one of claims 10 to 14 to produce a transformed plant;
- ii) selecting said transformed plant for occurrence of said isolated DNA molecule; and
- iv) growing said transformed plant in order to produce said protein of interest, wherein expression of said protein of interest is induced by the expression product of said isolated DNA.

56. The method of claim 55, wherein said transformed plant is transformed with a second vector comprising a nucleotide sequence encoding said protein of interest under the control of a regulatory element, said regulatory element induced by the expression product of said isolated DNA..

57. The method of claim 55, wherein said protein of interest is a native protein.

58. The method of any one of claims 55 or 56, wherein said protein of interest is selected from the group consisting of a pharmaceutically active protein, antibody, industrial enzyme, protein supplement, nutraceutical, storage protein, an enzyme involved in oil biosynthesis, animal feed, and animal feed supplement.

59. The isolated DNA molecule of claim of any one of claims 4 to 7, wherein said isolated DNA molecule encodes a protein that is at least 70% homologous with the amino acid defined by SEQ ID NO:2.

60. The isolated DNA molecule of claim of any one of claims 4 to 7, wherein said isolated DNA molecule encodes a protein that is at least 70% homologous with the amino acid defined by SEQ ID NO:4.

61. The isolated protein of claim 18, wherein said protein comprises from about 30 to about 541 amino acids of the sequence disclosed in SEQ ID NO:2

62. The isolated protein of claim 18, wherein said protein comprises from about from about 30 to about 561 amino acids of the sequence disclosed in SEQ ID NO: 4.

Figure 1




Figure 2

BNM3A	GTTCATCTCTCTCTTTAAGACCAAAACCTTTTCTCCTCTCTTCATGCATGAACCCCTA	60
BNM3B	-----TTCCTCTTTTACCTTTTACCAAGAACTCGTTAGATCAGCTCTGAACTCAA	120
BNM3A	ACTAAGTTCTCTCTCTTTTACCTTTTACCAAGAACTCGTTAGATCAGCTCTGAACTCAA	120
BNM3B	-----TTCCTCTTTTACCTTTTACCAAGAACTCGTTAGATCAGCTCTGAACTCAA	51
BNM3A	TGAATAATAACTGGTAGGCTTTTCTCTCTCTCTTATGAACAAAATCACCATCGTAAGG	180
BNM3B	TGAATAATAACTGGTAGGCTTTTCTCTCTCTCTTATGAACAAAATCACCATCGTAAGG	111
BNM3A	ACGCTCTACTCTTCCACCACCACAACCGTCGTAGATGTCGCCGGAGAGTACTGTTACGATC	240
BNM3B	ACGCTCTCTCTTCCACCACCACAACCGCGGTAGATGTCGCCGGAGAGTACTGTTACGATC	171
BNM3A	CGACCCTGCTCTCCGATGAGTCTTTCAGCGATCCAAACATCGTTTCTCTCTCTTTGGTG	300
BNM3B	CGACCCTGCTCTCCGATGAGTCTTTCAGCGATCCAAACATCGTTTCTCTCTCTTTGGTG	231
BNM3A	TCTGCTGATGCTTTTCCAGAGACAACAATAGTCACTCCCGAGATTGGGACATCAATG	360
BNM3B	TCTGCTGATGCTTTTCCAGAGACAACAATAGTCACTCCCGAGATTGGGACATCAATG	291
BNM3A	GTGTGTCATGCAATAACATCCACAACGATGAGCAAGTGGACCAAAGCTTGAGAAATTTTC	420
BNM3B	GTGTGTCATGTAATAACATCCACAATGATGAGCAAGTGGACCAAAGCTTGAGAAATTTTC	351
BNM3A	TTGGCCGCACCAACCGATTACAAACCAACGAAACGTGGAGATGGAAGTGGAGTG	480
BNM3B	TTGGCCGCACCAACCGATTACAAACCAACGAAACGTGGAGATGATGGAAGTG	411
BNM3A	GCTGTTATGGAGGAGAGACCGTGTGTGTGGCTCACTAGGACTTTGATGATAAAGACAT	540
BNM3B	GCTGTTATGGAGGAGAGACCGTGTGTGTGGCTCACTAGGACTTTGATGATAAAGACAT	471
BNM3A	GGCTGAGAAATCAACCGGTGGATAATGTTGATAATCAAGAAAATGGCAATGTCGCAAAAG	600
BNM3B	GGCTGAGAAATCAACCGGTGGATAATGTTGATAATCAAGAAAATGGCAATGTCGCAAAAG	531
BNM3A	GCTGTCTCTCTCAATGAATCATCTACTTCTGTGATAACAAACACAGCAGCAATAACA	660
BNM3B	GCTGTCTCTCTCAATGAATCATCTACTTCTGTGATAACAAACACAGCAGCAATAACA	591
BNM3A	ACCTGTTGTGCCCAAGGGAAGACTATTGATGATAGCGTTGAAGCTACACCGAAGAAAACCTA	720
BNM3B	ACCTGTTGTGCCCAAGGGAAGACTATTGATGATAGCGTTGAAGCTACACCGAAGAAAACCTA	651
BNM3A	TTGAGAGTTTGGACAGAGGACGTCTATATACCGCGGTGTTACAAGGCATCGGTGGACAG	780
BNM3B	TTGAGAGTTTGGACAGAGGACGTCTATATACCGCGGTGTTACAAGGCATCGGTGGACAG	711
BNM3A	GAGGATATGAGGCACATTTATGGGATATAGTTGTAAAGAGAAAGGCCAAACCGGCAAG	840
BNM3B	GAGGATATGAGGCACATTTATGGGATATAGTTGTAAAGAGAAAGGCCAAACCGGCAAG	771
BNM3A	GAGGACAAGTTTATTTGGGAGTTTATGACAAAGAAAGAACGAGCTAGGGCTTATGATT	900
BNM3B	GAGGACAAGTTTATTTGGGAGTTTATGACAAAGAAAGAACGAGCTAGGGCTTATGATT	831
BNM3A	TAGCCGCACTCAAGTATTGGGGAACCAACCACTACTACTAACTTCCCATGAGCGAATAG	960
BNM3B	TAGCCGCACTCAAGTATTGGGGAACCAACCACTACTACTAACTTCCCATGAGCGAATAG	891
BNM3A	AAAAAGAGGTAGAAGAGATGAAGCACATGACAAAGCAAGAGATGTTGCTCTACTGCGCA	1020
BNM3B	AAAAAGAGGTAGAAGAGATGAAGCACATGACAAAGCAAGAGATGTTGCTCTACTGCGCA	951
BNM3A	GGAAAAGTAGTGTTTCTCTCTGCGTGCATCATTTATCGTGAGTAACAAAGCATCAAC	1080
BNM3B	GGAAAAGTAGTGTTTCTCTCTGCGTGCATCATTTATCGTGAGTAACAAAGCATCAAC	1011

BNM3A	AACATGGAAGATGGCAAGCTAGGATAGGAAGAGTCGCCGGTAACAAAGACCTCTACTTGG	1140
BNM3B	AACATGGAAGATGGCAAGCTAGGATAGGAAGAGTCGCCGGTAACAAAGACCTCTACTTGG	1071
BNM3A	GAACCTTTGGCACACAAGAAGCTGAGAGGCATACGACATTGCGGCCATCAAAATCA	1200
BNM3B	GAACCTTTGGCACACAAGAAGCTGAGAGGCATACGACATTGCGGCCATCAAAATCA	1131
BNM3A	GAGGATTAAACCGCAGTGACTAATTCGACATGAACAGATACAAAGTTAAAGCAATCCTCG	1260
BNM3B	GAGGATTAAACCGCAGTGACTAATTCGACATGAACAGATACAAAGTTAAAGCAATCCTCG	1191
BNM3A	AAAGCCCTAGTCTTCCCTATGGTADCGCCGCAAAAGCTCTCAAGGAGGCTAACCGTCCGG	1320
BNM3B	AAAGCCCTAGTCTTCCCTATGGTADCGCCGCAAAAGCTCTCAAGGAGGCTAACCGTCCGG	1251
BNM3A	TTCCAAGTATGATGATGATCAGCAATAAAGCTTTCAGAGAGTGAGAAATAGTGCTAGCGGTT	1380
BNM3B	TTCCAAGTATGATGATGATCAGCAATAAAGCTTTCAGAGAGTGAGAAATAGTGCTAGCGGTT	1311
BNM3A	GGCAAAACGCTGCGGTTCAAGCATCATCAGGGAGTAGATTGAGCTTATTGCACCAACATC	1440
BNM3B	GGCAAAACGCTGCGGTTCAAGCATCATCAGGGAGTAGATTGAGCTTATTGCACCAACATC	1371
BNM3A	AAGAGAGGTACAATGGTTATTATTACAATGGAGGAACCTTGCTTCGGAGAGTGCAGGG	1500
BNM3B	AAGAGAGGTACAATGGTTATTATTACAATGGAGGAACCTTGCTTCGGAGAGTGCAGGG	1431
BNM3A	CTTGTTTCAAAACAGAGGATGATCAACACCAATTTCTTGAGCAACACGAGAGCCTCATGA	1560
BNM3B	CTTGTTTCAAAACAGAGGATGATCAACACCAATTTCTTGAGCAACACGAGAGCCTCATGA	1491
BNM3A	CTAATATCGATCATCAAAGTTCTGTTTCGGATGATTCGGTTACTGTTGTGGAAATGTTG	1620
BNM3B	CTAATATCGATCATCAAAGTTCTGTTTCAGATGATTCGGTTACTGTTGTGGAAATGTTG	1551
BNM3A	TTGGTTATGGTGGTTATCAAGGATTTGAGCCCGGTTAACTGCGATGCTACGCTGCTA	1680
BNM3B	TTGGTTATGGTGGTTATCAAGGATTTGAGCCCGGTTAACTGCGATGCTACGCTGCTA	1611
BNM3A	GTGAGTTTGATTATAACGCAAGAAACCATTTTACTTTGCTCAGCAGCAGACCCAGC	1740
BNM3B	GTGAGTTTGACTATAACGCAAGAAACCATTTTACTTTGCTCAGCAGCAGACCCAGC	1671
BNM3A	AGTCGCCAGGTGGAGATTTTCCCGCGGCAATGACGAATAATGTTGGCTCTAATATGTAATT	1800
BNM3B	ATTGCGCAGGAGGAGATTTTCCCGCGGCAATGACGAATAATGTTGGCTCTAATATGTAATT	1731
BNM3A	ACCATGGGGAAGGTGGTGGAGAAGTTGCTCCACATTTACAGTTTGGAAACGACAAATAGA	1860
BNM3B	ACCATGGGGAAGGTGGTGGAGAAGTTGCTCCACATTTACAGTTTGGAAACGACAAATAGA	1791
BNM3A	AAAAATAGTTAAAGATCTTAGTTATATGCGTTGTGTGTCGTTGTAACAGTGTGATAC	1920
BNM3B	AAAAATAGTTAAAGATCTTAGTTATATGCGTTGTGTGTCGTTGTAACAGTGTGATAC	1851
BNM3A	TTTGATTATGTTTTCCTCTCTCTTTCTCTTTCTCTGGTTAATTTCTAAGACTTTATT	1980
BNM3B	TTTGATTATGTTTTCCTCTCTCTTTCTCTTTCTCTGGTTAATTTCTAAGACTTTATT	1909
BNM3A	TTTAGTTTCCATTAGTTGGATAAATTTTCAGACT-----	2014
BNM3B	TTTAGTTTCCATTAGTTGGATAAATTTTCAGACT-----	1969
BNM3A	-----	
BNM3B	GTCTAATACAGAAAGTTTCAT	1992

Figure 3

```

BNM3A  MNNNLGFSLSPYEQNHHRKDVSSTTTTVDVAGEVCYDPTAASDESSAIQTSPFSPFG  60
BNM3B  MNNNLGFSLSPYEQNHHRKDVSSTTTTAVDVAGEVCYDPTAASDESSAIQTSPFSPFG
      *.....*

BNM3A  VVVDAFTRDNNSHSRWDINGCACNNHNDEQDGPKLENFLGRITTTIYNTNENVGDGSGS  120
BNM3B  VVLDAFTRDNNSHSRWDINGACNNHNDEQDGPKLENFLGRITTTIYNTNENVGDIDGS
      *.....*

BNM3A  GCYGGGDDGGGSLGLSMIKTWLRNPVDNVDNQENGNAAGKLSLSMNSSTSCDNNNDNN  180
BNM3B  GCYGGGDDGGGSLGLSMIKTWLRNPVDNVDNQENGNAAGKLSLSMNSSTSCDNNNYSSN
      *.....*

      repeat 1
BNM3A  NLVAQGGKTIDDSVEATPKKTIIESFGQRTSIYRGVTRHRNTGRYEAHLWDNSCKREGQTRK  240
BNM3B  NLVAQGGKTIDDSVEATPKKTIIESFGQRTSIYRGVTRHRNTGRYEAHLWDNSCKREGQTRK
      *.....*

BNM3A  GRQVYLGGYDREEKAARAYDLAALKYWGTTTTNPFMSEYEKEVEEMKHMTRQEYVASLR  300
BNM3B  GRQVYLGGYDREEKAARAYDLAALKYWGTTTTNPFMSEYEKEIEEMKHMTRQEYVASLR
      *.....*

      repeat 2
BNM3A  RKSSGFSRGASIIYRGVTRHHQHGRWQARIGRVAGNKDLYLGTFTGTQEEAAEAYDIAAIFK  360
BNM3B  RKSSGFSRGASIIYRGVTRHHQHGRWQARIGRVAGNKDLYLGTFTGTQEEAAEAYDIAAIFK
      *.....*

BNM3A  RGLTAVTNFDMNRYYNKAILESPLIGSAAKRLKEANRPVPSMMHISNNVSESENASG  420
BNM3B  RGLTAVTNFDMNRYYNKAILESPLIGSAAKRLKEANRPVPSMMHISNNVSESENASG
      *.....*

BNM3A  WQNAAVQHHCQVDLSLLHQHQERYNGYYINGNLSSSARACFKQEDDQHFLSNTQSLM  480
BNM3B  WQNAAVQHHCQVDLSLLHQHQERYNGYYINGNLSSSARACFKQEDDQHFLSNTQSLM
      *.....*

BNM3A  TNIDHQSSVSDSVTVCGNVVGYGGYQGFAPVNCDAYAASEFDYNARNHYFAQQQQTQ  540
BNM3B  TNIDHQSSVSDSVTVCGNVVGYGGYQGFAPVNCDAYAASEFDYNARNHYFAQQQQTQ
      *.....*

BNM3A  QSPGGDFPAAMTNNVGSNNYYHGGEGGEVAPFTVWNDN  579
BNM3B  HSPGGDFPAAMTNNVGSNNYYHGGEGGEVAPFTVWNDN
      *.....*

```

Figure 4




Figure 5

AP2 DOMAIN REPEAT 1

BNM3 TSIYRGVTRHRWTRGYEAHLWDNSCKREGQTRKQVYLGVDKKEKAAAYDLAALKYWGTTTTFNPFMSEYEK
 ANT T*Q******F*K*HS*****M*****PS*H**SAEN*Q*I
 ZM T******E*V*R*S*****F*P*P*****QV*N**L
 GL15 S*Q*+++FY*R**W*S*I*+---C-----*K*****F*TAQA*****Q*I*PR*LNADI*TLDD*KD*M
 AP2 S*Q*+++FY*R**W*S*I*+---C-----*K*****F*TAHA*****R*I*PR*VEADI*NIDD*DDDL

LINKER

BNM3 EEMKHMTRQEYVASLRKSSGFSRG
 ANT *D*N*****H*****
 ZM ***S*****EI*****
 GL15 KK*DLKKE*F*LV**QGA**V*
 AP2 KQ*TNL*KE*F*HV**Q*T**P**

AP2 DOMAIN REPEAT 2

BNM3 ASIYGVTRHHQHGRWQARIGRVAGNKDYLGTGFTGTEEAAYDIAAIKFRGLTAVTNFDMNRYNVKA
 ANT *****S*****S*****S*****TN*****IT**D*DR
 ZM *****E*****V*****N**L*S**D*ES
 GL15 S*RF*+*+*Q*RC*K*E*****QLM*K*YV***LYD*ET***Q**K***CY*KE*****AQS*DKEL
 AP2 S*K*+*+*+*L*KC*+*E**M*QFL*K*YV***L*D*EV***R**K*****CN*KD*****PSI*DBEL

Figure 6




Figure 7






Figure 8





European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 99 20 1745

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	NAKAMURA Y.: "AC AB025629" EMBL DATABASE, 9 April 1999 (1999-04-09), XP002120887 Heidelberg * the whole document *	1,2	C12N15/29 C12N15/82 C12N5/10 C07K14/415 A01H5/00 A01H5/10
X	RIEGER M. ET AL.: "AC AL049862" EMBL DATABASE, 13 May 1999 (1999-05-13), XP002120888 Heidelberg * the whole document *	28	
X,D	ELLIOTT R. ET AL.: "AINTEGUMENTA, an APETALA2-like gene of arabidopsis with pleiotropic roles in ovule development and floral organ growth" THE PLANT CELL, vol. 8, 1996, pages 155-168, XP002120889 * see esp. p.161;165 *	54	
A,D	WO 97 43427 A (CIBA GEIGY AG ;VRIES SAPE CORNELIS DE (NL); SCHMIDT EDUARD DANIEL) 20 November 1997 (1997-11-20) * the whole document *	1-62	TECHNICAL FIELDS SEARCHED (Int.Cl.7) C12N C07K
A	WO 98 37184 A (UNIV CALIFORNIA) 27 August 1998 (1998-08-27) * see esp. example 2 *	1-62	
A	WO 98 07842 A (JOFUKU K DIANE ;OKAMURO JACK K (US); UNIV CALIFORNIA (US)) 26 February 1998 (1998-02-26) * the whole document *	1-62	
-/--			
This present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 29 October 1999	Examiner Kania, T
CATEGORY OF CITED DOCUMENTS		I : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons A : technological background O : non-written disclosure P : intermediate document	
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		& : member of the same patent family, corresponding document	

EPO FORM 1503 (3.82) (PUB.01)

European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 99 20 1745

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
A,D	OGAS J. ET AL.: "Cellular differentiation regulated by gibberellin in the Arabidopsis thaliana pickle mutant" SCIENCE, vol. 277, July 1997 (1997-07), pages 91-94, XP002120890 * the whole document *	1-62	TECHNICAL FIELDS SEARCHED (Int.Cl.7)
A,D	BOUTILIER, K. A. ET AL.: "Expression of the BnmNAP subfamily of napin genes coincides with the induction of Brassica microspore embryogenesis" PLANT MOLECULAR BIOLOGY, (1994) VOL. 26, NO. 6, PP. 1711-1723. 48 REF. ISSN: 0167-4412, XP002120891 * the whole document *	1-62	
A	CUSTERS, JAN B.M. ET AL.: "Regulation of the inductive phase of microspore embryogenesis in Brassica napus" ACTA HORTIC. (1996), 407 (INTERNATIONAL SYMPOSIUM ON BRASSICAS, NINTH CRUCIFER GENETICS WORKSHOP, 1994), 209-217, XP002120892 * see the whole document; esp. table 3 clone 13 *	1-62	
A	CHAUDHURY A. ET AL.: "Ovule and embryo development, apomixis and fertilization" CURRENT OPINION IN PLANT BIOLOGY, 'Online!' vol. 1, 1998, pages 26-31, XP002120893 Retrieved from the Internet: <URL:http://biomednet.com/eleceref/1369526600100026> 'retrieved on 1999-10-27! * see esp. p.29 *	1-62	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 29 October 1999	Examiner Kania, T
CATEGORY OF CITED DOCUMENTS		T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons A: technological background O: non-written disclosure P: intermediate document &: member of the same patent family, corresponding document	

EPO FORM 1503 (02/82) (P4/C31)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 99 20 1745

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

29-10-1999

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9743427	A	20-11-1997	AU	2953997 A	05-12-1997
			CA	2254839 A	20-11-1997
			CN	1218510 A	02-06-1999
			EP	0915984 A	19-05-1999
			HU	9901477 A	28-09-1999
			PL	329872 A	12-04-1999
WO 9837184	A	27-08-1998	AU	6328398 A	09-09-1998
WO 9807842	A	26-02-1998	AU	3986797 A	06-03-1998
			PL	331771 A	02-08-1999

EPO FORM P/2002

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82